

Data Papers

Ecology, 91(3), 2010, p. 930
© 2010 by the Ecological Society of America

The Jena Experiment: six years of data from a grassland biodiversity experiment

Ecological Archives E091-066

ALEXANDRA WEIGELT,^{1,6} ELISABETH MARQUARD,^{2,3} VICKY M. TEMPERTON,^{3,7} CHRISTIANE ROSCHER,³ CHRISTOPH SCHERBER,⁴ PETER N. MWANGI,^{1,8} STEFANIE VON FELTEN,² NINA BUCHMANN,⁵ BERNHARD SCHMID,² ERNST-DETLEF SCHULZE,³ AND WOLFGANG W. WEISSE¹

¹Institute of Ecology, University of Jena, Dornburger Strasse 159, D-07743 Jena, Germany

²Institute of Environmental Sciences, University of Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

³Max Planck Institute for Biogeochemistry, Hans-Knoll Strasse 10, D-07745 Jena, Germany

⁴Department of Crop Sciences, University of Göttingen, Waldweg 26, D-37073 Göttingen, Germany

⁵ETH Zürich, Universitätstr. 2, CH-8092 Zürich, Switzerland

Abstract. This data set contains species-specific biomass and cover data as well as community leaf area index (LAI) and height from a large grassland biodiversity experiment (Jena Experiment). In this experiment, 82 grassland plots of 20 × 20 m were established from a pool of 60 species belonging to four functional groups (grasses, legumes, and tall and small herbs). In May 2002, varying numbers of plant species were sown into the plots to create a gradient of plant species richness (1, 2, 4, 8, 16, and 60 species) and functional richness (1, 2, 3, and 4 functional groups). Plots were maintained by biannual weeding and mowing. The data set encompasses the 2002–2008 May and August biomass harvests from 3–4 subplots of 0.2 × 0.5 m per experimental plot sorted to species. Moreover, plant species and community cover estimated in an approximately 9-m² subplot per plot are included in the data set. Each biomass harvest was accompanied by measurements of vegetation height and LAI per plot. Analyses of the community biomass data have identified species richness as well as functional group composition as important drivers of a positive biodiversity–productivity relationship.

The data set can be used to study a variety of questions about how plant community composition and structure respond to changes in species richness and functional diversity over time. Sampling is ongoing, and new data will be added.

Key words: biodiversity experiment; functional composition; height; LAI (leaf area index); plant community; species biomass; species cover.

The complete data sets corresponding to abstracts published in the Data Papers section of the journal are published electronically in *Ecological Archives* at <<http://esapubs.org/archive>>. (The accession number for each Data Paper is given directly beneath the title.)

Manuscript received 24 May 2009; revised 10 September 2009; accepted 22 September 2009. Corresponding Editor: W. K. Michener.

⁶E-mail: alexandra.weigelt@uni-jena.de

⁷Present address: Phytosphere (ICG-3), Forschungszentrum Jülich GmbH, D-52425 Jülich, Germany.

⁸Present address: Department of Botany, Jomo Kenyatta University of Agriculture and Technology, P.O. Box 62000-00200 Nairobi, Kenya.