

Labile carbon retention compensates for CO₂ released by priming in forest soils

NA QIAO*†¹, DOUGLAS SCHAEFER*¹, EVGENIA BLAGODATSKAYA‡§, XIAOMING ZOU*¶, XINGLIANG XU*‡ and YAKOV KUZYAKOV‡§||

*Key Laboratory of Tropical Forest Ecology, Chinese Academy of Sciences, Xishuangbanna Tropical Botanical Garden, Menglun, Mengla, Yunnan 666303, China, †Graduate School of the Chinese Academy of Sciences, 19A Yuquan Road, Beijing 100049, China, ‡Key Laboratory of Ecosystem Network Observation and Modeling, Chinese Academy of Sciences (CAS), Institute of Geographic Sciences and Natural Resources Research, 11A Datun Road, Chaoyang District, Beijing 100101, China, §Department of Soil Science of Temperate Ecosystems, University of Göttingen, Büsgenweg 2, Göttingen 37077, Germany, ¶Institute for Tropical Ecosystem Studies, University of Puerto Rico, P.O. Box 21910, San Juan, Puerto Rico 00931-1910, USA, ||Department of Agricultural Soil Science, University of Göttingen, Büsgenweg 2, Göttingen 37077, Germany

Abstract

Increase of belowground C allocation by plants under global warming or elevated CO₂ may promote decomposition of soil organic carbon (SOC) by priming and strongly affects SOC dynamics. The specific effects by priming of SOC depend on the amount and frequency of C inputs. Most previous priming studies have investigated single C additions, but they are not very representative for litterfall and root exudation in many terrestrial ecosystems. We evaluated effects of ¹³C-labeled glucose added to soil in three temporal patterns: single, repeated, and continuous on dynamics of CO₂ and priming of SOC decomposition over 6 months. Total and ¹³C labeled CO₂ were monitored to analyze priming dynamics and net C balance between SOC loss caused by priming and the retention of added glucose-C. Cumulative priming ranged from 1.3 to 5.5 mg C g⁻¹ SOC in the subtropical, and from -0.6 to 5.5 mg C g⁻¹ SOC in the tropical soils. Single addition induced more priming than repeated and continuous inputs. Therefore, single additions of high substrate amounts may overestimate priming effects over the short term. The amount of added glucose C remaining in soil after 6 months (subtropical: 8.1–11.2 mg C g⁻¹ SOC or 41–56% of added glucose; tropical: 8.7–15.0 mg C g⁻¹ SOC or 43–75% of glucose) was substantially higher than the net C loss due to SOC decomposition including priming effect. This overcompensation of C losses was highest with continuous inputs and lowest with single inputs. Therefore, raised labile organic C input to soils by higher plant productivity will increase SOC content even though priming accelerates decomposition of native SOC. Consequently, higher continuous input of C belowground by plants under warming or elevated CO₂ can increase C stocks in soil despite accelerated C cycling by priming in soils.

Keywords: ¹³C, addition frequency, carbon balance, glucose, litter decomposition, priming effect, root exudates, soil organic matter stability, subtropical forest, tropical forest

Received 11 October 2013 and accepted 17 October 2013

Introduction

The global pool of soil organic carbon (SOC) has been estimated to exceed 3300 Pg C (Tarnocai *et al.*, 2009; German *et al.*, 2011), four times larger than the atmospheric CO₂-C pool and five times greater than the C pool in vegetation. Therefore, even small changes in SOC would have large effects on atmospheric CO₂ and have potential feedbacks to climate (Kirschbaum, 2004; Heimann & Reichstein, 2008), emphasizing the need to understand SOC dynamics. In the past 20 years, an increasing number of studies have shown that inputs of

labile organic carbon (LOC) greatly enhance native SOC decomposition (Fontaine *et al.*, 2004a, 2007; van Hees *et al.*, 2005; Blagodatskaya *et al.*, 2007); known as 'priming effects' (Kuzyakov *et al.*, 2000). Therefore, priming effects have the potential to change SOC dynamics.

Root exudation and litter decomposition supply LOC to soil microorganisms and are key processes of C cycling in terrestrial ecosystems (Cheng *et al.*, 2013). They play important roles in maintaining soil functions and are strongly affected by environmental changes. Numerous studies have observed that both warming and elevated CO₂ can produce more litterfall and root exudates via increasing ecosystem net primary production (NPP, e.g., Pendall *et al.*, 2004; Finzi *et al.*, 2007; Jackson *et al.*, 2009; Wu *et al.*, 2011). Increased litter

Correspondence: Xingliang Xu, tel. +86 10 64889813, fax +86 10 64889813, e-mail: xuxingl@hotmail.com

¹These authors wish to be considered joint first authors.

decomposition induced by warming has also been demonstrated in a variety of ecosystems (Liski *et al.*, 2003). If C stored in soils is released to the atmosphere by SOC priming, there would be a positive feedback on climate. Conversely, if increases of plant-derived C inputs to soils exceed temperature-driven increases in litter decomposition, the feedback would be negative (Davidson & Janssens, 2006). A full understanding of priming effects induced by LOC inputs will be critical to clarify effects of warming and elevated CO₂ on soil C cycling and its feedback to global change (Cheng *et al.*, 2013).

Most previous priming experiments focused on single LOC additions, and showed that priming can accelerate decomposition of SOC (Blagodatskaya & Kuzyakov, 2008). However, single additions may not be representative of natural C inputs in terrestrial ecosystems. For example, in forests lacking strong seasonality, root exudates and surface litter are produced throughout the year. Even in strongly seasonal climates, dissolved organic C (DOC) input from root exudation and litter decomposition occurs continuously during the growing season. Root exudation also varies at the spatial scales of microorganisms. As a growing root tip extends through a zone of soil, microorganisms can experience a 'pulse' of exudation. Such variations in time and space could be simulated by repeated additions. Evidence suggests that a small change in the magnitude of priming effects has the potential to outweigh the effects of global change drivers on basal soil organic matter (SOM) decomposition in many ecosystems (Cheng *et al.*, 2013). Therefore, to clarify whether single, repeated, and continuous LOC inputs cause different amounts of priming is one of key questions for understanding the significance of priming and its occurrence. If single additions induce more priming than repeated or continuous additions, the former could overestimate priming occurring in natural ecosystems. So far no studies have compared priming effects induced by single, repeated, and continuous inputs in the same experiment, although a few studies have examined priming effects induced by repeated (Hamer & Marschner, 2005; Chigineva *et al.*, 2009) or continuous additions of LOC (Kuzyakov *et al.*, 2007; Paterson & Sim, 2013).

Priming increases SOC decomposition (e.g., Fontaine *et al.*, 2004a, 2007; Sayer *et al.*, 2011), but a fraction of the added organics can remain in the soil and compensate for the SOC loss caused by priming, potentially leading to a net C increase in soil (Fontaine *et al.*, 2004b; Ohm *et al.*, 2007). To understand effects of priming on SOC dynamics, the net balance between primed SOC loss and LOC retention must be known. However, most priming studies have emphasized SOC losses induced by priming and have not reported net SOC balances between primed C and the gain from added

LOC (Kuzyakov, 2010). Therefore, despite the overall importance of priming, net effects of LOC inputs on SOC dynamics and stabilization remain unclear.

Tropical and subtropical forests contain more than 20% of the global SOC (Tarnocai *et al.*, 2009), and are important sources of CO₂ return to the atmosphere (Raich *et al.*, 2002), indicating crucial roles of these forests in the global C cycle (Reich, 2011; Wood *et al.*, 2012; Cox *et al.*, 2013). Elevated atmospheric CO₂ has increased photosynthesis and productivity of tropical forest (Ziska *et al.*, 1991; Lewis *et al.*, 2009; Cernusak *et al.*, 2013) and understory (Würth *et al.*, 1998) species. More than 50 years of data also suggest that elevated atmospheric CO₂ is a driver for increased tree density in South African savannas (Buitenwerf *et al.*, 2012). Although some studies suggest that tropical forests are not sensitive to warming, litter production, tree growth, and belowground C allocation all increase significantly with mean annual temperature (Raich *et al.*, 2006). This indicates that both warming and elevated CO₂ have potential to increase NPP and litter production, and warming can accelerate litter decomposition in subtropical and tropical forests (Liski *et al.*, 2003). Free-air CO₂ enrichment studies have so far been extratropical, and most of them found net SOC increases (Hoosbeek *et al.*, 2004; Drigo *et al.*, 2008, 2013; Hungate *et al.*, 2009; Liu *et al.*, 2009; Phillips *et al.*, 2009, 2011; Dieleman *et al.*, 2010; Drake *et al.*, 2011). Modeling by Hickler *et al.* (2008) suggest that tropical forests may be even more sensitive to CO₂ enrichment than temperate ones.

Subtropical and tropical wet forests are dominated by evergreen broad-leaved tree species, and root exudation and litter decomposition can occur throughout the year. Within such forests, there are wide ranges of vegetations, climates, and soils. We chose two very different forests (2.8% SOC in the tropical and 13.8% SOC in the subtropical) to explore priming induced by LOC inputs. Total rhizodeposition in forests has rarely been quantified *in situ*, but available evidence suggests that root exudates can constitute 10% or more of NPP (Chapin *et al.*, 2012). According to de Graaff *et al.* (2010), the expected range for root exudation varies from 50 to 1500 µg C g⁻¹ soil per day (Trofymow *et al.*, 1987; Meharg & Kilham, 1991; Jones & Darrah, 1993; Cheng, 1996). In addition, litter decomposition releases DOC into soils. Elevated CO₂ can increase DOC input to soils, by increasing both root exudation and litter decomposability (Phillips *et al.*, 2008, 2009). Accurate evaluations of priming caused by LOC inputs in these forests will improve our understanding of C cycling at low latitudes.

There are large areas of tropical and subtropical forests in southern China (Wu, 1980). Although litter production in our tropical forest is double that in the subtropical (Schaefer *et al.*, 2009; Tang *et al.*, 2010), litter

decomposes more rapidly in the former (Liu *et al.*, 2000; Tang *et al.*, 2010). Slower litter decomposition at lower temperatures contributes to fivefold higher SOC content in subtropical soils (Ailaoshan) compared with tropical soils (Xishuangbanna). This tropical forest has $900 \text{ g C m}^{-2} \text{ yr}^{-1}$ of NPP (Tan *et al.*, 2010), with $1000 \text{ g C m}^{-2} \text{ yr}^{-1}$ in the subtropical (Tan *et al.*, 2011). Assuming that 10% of NPP becomes root exudates (Chapin *et al.*, 2012), the current rates are on the order of $100 \text{ g C m}^{-2} \text{ yr}^{-1}$ in these forests.

Although a few studies have examined priming at low latitudes (e.g., Fontaine *et al.*, 2004b; Schaefer *et al.*, 2009; Sayer *et al.*, 2011; Leff *et al.*, 2012), many details remain unclear. To clarify whether single vs. more frequent glucose inputs produce different priming effects, we made single (all at the beginning), repeated (once per month), and continuous (once per week) additions. The same total amount of glucose was added to each treatment over the study period. Lower C/N in the tropical than in the subtropical soil suggests faster SOC turnover in the tropical soil (Krull & Skjemstad, 2003; Rumpel & Kögel-Knabner, 2011). We therefore hypothesized that greater SOC priming would occur in the tropical soil. Because microbial responses might persist longer with continuous inputs than after a single addition, we also hypothesized that continuous additions would result in stronger priming than a single addition of the same total amount of glucose. Our first glucose additions differed in amount between treatments, so we could examine the linearity of initial microbial responses. As priming accelerates SOC decomposition (e.g., Fontaine *et al.*, 2004a, 2007; Sayer *et al.*, 2011), we hypothesized that the net C balances between primed C and retention of added glucose-C could be negative in both forest soils. To test these hypotheses, we conducted 170-day incubations of tropical (Xishuangbanna) and subtropical (Ailaoshan) forest soils.

Materials and methods

Soils

After removing the litter layer, we sampled mineral soil from the top 10 cm in subtropical and tropical forests in July 2009. The subtropical forest was in the Ailao Mountains

Nature Reserve (24°32' N, 101°01' E, 2476 m asl) in the Yunnan Province of southwestern China. This forest is dominated by the evergreen broad-leaved species *Lithocarpus chintungensis*, *Rhododendron leptothrium*, *Vaccinium duclouxii*, *Lithocarpus xylocarpus*, *Castanopsis wattii*, *Schima noronhae*, *Hartia sinensis*, and *Manglietia insignis* (Li, 1983). Mean annual air temperature is 11 °C and annual precipitation averaged 1950 mm from 1996 to 2006. The monsoon climate causes a wet season from May to October followed by a dry season from November to April (Zhang, 1983). The soil is a loamy Lixisol (World Reference Base for Soil Resources, 2006) or Ustalf Alfisol (by USDA Soil Taxonomy) derived from weakly metamorphosed marine sediments.

The tropical forest site was in the Xishuangbanna Tropical Rainforest Ecosystem Station (21°54' N, 101°16' E, 560 m asl) in Menglun, Xishuangbanna, also in Yunnan Province. Annual mean temperature and precipitation are 21.6 °C and 1480 mm. About 85% of the rain falls from May to October. The tree layer is dominated by *Pometia tomentosa*, *Gironniera subaequalis*, *Chisocheton siamensis*, *Barringtonia macrostachya*, and *Pseuderanthemum latifolium*. The soil is a Ferrasol (World Reference Base for Soil Resources, 2006) or Oxisols (by USDA Soil Taxonomy) derived from marine sandstone (Table 1).

Experimental design

Thirty grams of air-dried, 2-mm-sieved, root-picked soil were added to 250-ml Schott bottles, and adjusted to 60% of the water-holding capacity (WHC). All soils were pre-incubated at 20 °C for 7 days. Thereafter, water or ^{13}C -uniformly-labeled glucose (5.97 atom% ^{13}C) solution was added evenly dropwise to the soil surface using a pipette to obtain uniform distribution. These additions raised soils to 70% of WHC and they were maintained at this level throughout the experiment. One gram of CaCl_2 was added to small cups placed in the incubation bottles to absorb water vapor and to prevent soil moisture increases from subsequent water additions. The CaCl_2 was replaced weekly to maintain its water-absorptive capacity.

Assuming that 10% of NPP becomes root exudates (Chapin *et al.*, 2012), the current rates are on the order of $100 \text{ g C m}^{-2} \text{ yr}^{-1}$ in these forests. In addition to root exudation, litter decomposition also can release LOC into soils. Decomposing litter can release 3% of its C as DOC (Don & Kalbitz, 2005), which would correspond to some $10 \text{ g C m}^{-2} \text{ yr}^{-1}$ in these forests. Allison *et al.* (2010) used an annual DOC flux of 44 g C m^{-2} to the top 1 cm of soil surface in their model. Elevated CO_2 also can increase LOC input to soils through increasing root exudates and litter decomposition (Phillips *et al.*, 2009). Hoosbeek & Scarascia-Mugnozza

Table 1 Properties of the surface 10 cm mineral soils in a subtropical forest at Ailaoshan and in a tropical forest at Xishuangbanna in Yunnan Province, China. Means \pm 1 SE, $n = 4$

Soil	Organic carbon (% dry wt.)	Total nitrogen (% dry wt.)	Microbial biomass carbon (mg g^{-1} dw soil)	C/N	WHC (%)
Subtropical forest	13.84 \pm 0.20	0.88 \pm 0.01	2.50 \pm 0.20	15.64 \pm 0.13	95.1 \pm 0.2
Tropical forest	2.78 \pm 0.01	0.28 \pm 0.00	0.52 \pm 0.10	9.98 \pm 0.02	57.5 \pm 3.5

(2009) summarized several FACE studies (typically doubling CO₂) as increasing SOC by 50 g C m⁻² yr⁻¹ or more (mostly from increased litterfall). Phillips *et al.* (2011) estimated total root exudation in a pine forest as 10 g C m⁻² yr⁻¹, and we speculate that doubled CO₂ could increase exudation by a similar amount. Based on these soils' %SOC, bulk density, 10 cm soil depth, and the glucose-C added (g⁻¹ SOC) in this experiment; our additions correspond to 110 and 280 g C m⁻² yr⁻¹ at the field scale to the tropical and subtropical soils. Our glucose-C additions were thus of the magnitude expected in these soils for current annual root exudation (de Graaff *et al.*, 2010) and DOC from litter decomposition. These additions were also comparable to microbial biomass-C (MBC), and so could change microbial activity but not be sufficient to induce obvious microbial growth (Blagodatskaya & Kuzyakov, 2008). To maintain comparability in these two different soils, total glucose-C additions were equal to 2% of the SOC in each soil (Table 1).

Single-addition treatments received all the glucose-C at the start of the experiment (after one week of pre-incubation) and all subsequent weekly additions were water only. Repeated addition treatments received 1/6 of the total glucose-C at monthly intervals with the intervening weekly additions being water only. Continuous-addition treatments received 1/24 of the total glucose-C each week. Controls received water-only additions each week. Each treatment had four replicates. The first glucose additions to each soil treatment varied in amount between treatments, but on an SOC-specific basis they were the same in corresponding treatments for both soils. This allowed us to examine whether different glucose additions saturated microbial utilization.

All incubations were conducted at 20 °C. Five milliliters of 1 M NaOH were placed in small cups in each incubation bottle to trap CO₂ and were replaced at the end of each week. CO₂ samples were trapped from each incubation bottle and analyzed for CO₂ and δ¹³C.

CO₂ analysis

To measure CO₂ absorbed in NaOH, 4 ml of 0.5 M SrCl₂ was added to precipitate carbonate. Unreacted NaOH was titrated with 0.2 M HCl against the phenolphthalein endpoint (Zibilske, 1994). Precipitated SrCO₃ was centrifuged three times at 1200 g for 10 min followed each time by rinsing with degassed water. The SrCO₃ was then dried at 105 °C and weighed into tin capsules to analyze for total C and ¹³C/¹²C ratios by continuous flow gas isotope ratio mass spectrometry (MAT253; Finnigan MAT, Bremen, Germany), coupled by ConFlo III device (Finnigan MAT) to a elemental analyzer (EA 1112; CE Instruments, Milan, Italy).

Microbial biomass and dissolved organic carbon

After the incubations, MBC was measured by chloroform fumigation/extraction (modified after Vance *et al.*, 1987). Briefly, 5 g fresh soil was extracted with 20 ml of 0.05 M K₂SO₄. An additional 5 g soil was fumigated with ethanol-free chloroform for 24 h, and then was extracted again in the same

manner. Total organic C concentrations in the K₂SO₄ extracts were measured with a Dimatec-100 TOC/TIC analyzer (Dimatec Analysentechnik GmbH, Essen, Germany). Total organic C concentrations in the K₂SO₄ extracts from nonfumigated soils were defined as DOC. Aliquots of the K₂SO₄ extracts were pipetted directly into tin capsules and dried at 60 °C to measure δ¹³C in the microbial biomass by mass spectrometry (Brant *et al.*, 2006).

Calculations

Total CO₂-C in trapped CO₂ was analyzed weekly and their values were reported on a specific basis (i.e. per unit of SOC). For each soil and treatment, CO₂ fluxes from the four replicates of were summed through the experiment, and cumulative mean values and standard deviations (SDs) were calculated for each week. Flux-weighted δ values were based on weekly values and CO₂ fluxes.

The end-member mixing model used to calculate fractions of CO₂-C derived from SOC (C_{SOC}) and from added glucose (C_{glucose}) was from Phillips & Gregg (2001) and Phillips *et al.* (2005), <http://www.epa.gov/wed/pages/models.htm>. This model allows variability from mass-spectrometric measurements to be combined with that from CO₂ flux measurements.

Based on the fractions of glucose-derived CO₂ and their SDs obtained from the model by Phillips & Gregg (2001), primed C was calculated as follows:

$$\text{Primed C} = C_{\text{total}} - C_{\text{glucose}} - C_{\text{water only}}$$

where C_{total} is total C-CO₂ from the glucose-treated soil, C_{glucose} is C-CO₂ derived from added glucose, and C_{water only} is total C-CO₂ from the soil adding only water.

Glucose release (%) was C_{glucose} divided by total glucose added, and net C balance was calculated as the difference between primed C and retention of added glucose-C.

Standard deviations of cumulative CO₂ fluxes, glucose release (%), priming, and net C balances were calculated according to the following equations described by Ku (1966).

$$\sigma_{x+y} = \sqrt{(\sigma_x)^2 + (\sigma_y)^2}$$

where σ_{x+y} is SD of the combined flux and σ_x and σ_y are SDs of the individual fluxes.

$$\sigma_{xy} = \bar{x} * \bar{y} * \sqrt{\frac{(\sigma_x)^2}{\bar{x}^2} + \frac{(\sigma_y)^2}{\bar{y}^2}}$$

where σ_{xy} is SD of the combined flux, σ_x and σ_y are SDs of the CO₂ flux and SOC fractions, and \bar{x} and \bar{y} are their respective mean values.

At the end of the incubations, MBC content was calculated as the difference between the total C in fumigated and nonfumigated soils, divided by a k_{EC} factor of 0.45 (Wu *et al.*, 1993). The δ¹³C of DOC was measured from the K₂SO₄ extracts from nonfumigated soils. The δ¹³C of the microbial biomass was calculated as the difference of ¹³C between fumigated and nonfumigated samples divided by the difference of the C amount between fumigated and nonfumigated samples (Blagodatskaya *et al.*, 2011). With these δ¹³C values

of MBC and DOC, the Phillips & Gregg (2001) end-member mixing model was also used to estimate the fraction of added glucose incorporated to the DOC and MBC pools.

Statistics

Tukey's HSD was used to test the effects of the pattern of glucose addition on primed SOC, retained glucose, and net C balances among treatments and between soils. Repeated measures ANOVAS (Pallant, 2007) over time were performed with SPSS 21.0 (SPSS Inc., Chicago, IL, USA) to evaluate the effects of soil type and the pattern of glucose addition on CO₂ flux rates. Standardized major axis regressions were used to test for differences in the CO₂ slopes and intercepts against substrate additions to the subtropical and tropical forest soils using R 3.0.1 with the SMATR package (Falster *et al.*, 2006; Warton *et al.*, 2006; R Development Core Team, 2011). All differences were tested for significance at $P = 0.05$.

Results

CO₂ efflux responses to glucose additions

Throughout incubations of the water-only controls, specific CO₂ efflux rates from the tropical soil ($169 \pm 8 \mu\text{g C g}^{-1} \text{SOC d}^{-1}$) were significantly higher than from the subtropical soil ($94 \pm 4 \mu\text{g C g}^{-1} \text{SOC d}^{-1}$; Fig. 1a,b). The CO₂ effluxes depended on glucose input patterns (Fig. 1a,b) with increases after each glucose addition. Repeated-measures ANOVA indicated that soil type and glucose addition patterns significantly affected CO₂ efflux rates (Table 2).

Table 2 Results of repeated-measures ANOVA over time to evaluate effects of soil type and the frequency of LOC addition on CO₂ flux rates. Bold numbers indicate significant differences at $P < 0.05$

Source of variation	df	MS	F	P
Between subjects				
Intercept	1.00	22946752	9136.34	0.000
Soil type	1.00	869964	346.38	0.000
Adding frequency	3.00	469036	62.25	0.000
Soil type × Adding frequency	3.00	8193	1.09	0.373
Error	24.00	60278		
Within subjects				
Time	6.94	3451797	75.71	0.000
Time × Soil type	6.94	338534	7.43	0.000
Time × Adding frequency	20.82	7358948	53.80	0.000
Time × Soil type × Adding frequency	20.82	1105378	8.08	0.000
Errors	166.56	1094247		

Glucose release as CO₂

Expressed as percent of the amount added, release of glucose CO₂ differed among treatments (Fig. 1c,d; Table S1, Tukey's HSD test, $P < 0.05$). In the tropical soil, continuous additions released less glucose than repeated additions during the incubation, but finally converged with the single additions ($28.8 \pm 0.7\%$). More glucose-CO₂ was respired from the repeated addition ($43.1 \pm 0.02\%$) to the tropical soil than from the other two treatments at the end of incubations (Fig. 1c, Table S1). In the subtropical soil, the pattern of glucose respiration was similar to that in the tropical soil, but their final fractions were significantly different among the three treatments (repeated $41.9 \pm 0.9\% >$ continuous $37.5 \pm 1.4\% >$ single $32.2 \pm 1.0\%$; Fig. 1d; Table S1). By the end of the incubations, single and continuous additions to the subtropical soil increased glucose-CO₂ release by 12% and 35% compared with the tropical soil. Compared with the tropical soil, repeated additions to the subtropical soil decreased by 3% glucose-CO₂ loss (tropical vs. subtropical: $43.1 \pm 0.02\%$ vs. $41.1 \pm 0.9\%$, $P < 0.05$; Fig. 1c,d; Table S1).

Dynamics of cumulative priming effects

Soil type and glucose-addition patterns significantly affected priming intensity and dynamics (Fig. 1, Table S2, Tukey's HSD test, $P < 0.05$). In the tropical soil, cumulative priming caused by single addition was positive and significantly greater than with repeated and continuous additions. The repeated and continuous additions produced similar priming and the values fluctuated around zero throughout the incubations (Fig. 1e; Table S2). In the subtropical soil, single additions also caused more priming than repeated and continuous additions throughout the incubations. However, the dynamics of priming were distinctly different between soils. Single addition quickly reached maximal priming in the tropical soil, with only minor changes later. In the subtropical soil, the priming induced by single addition increased gradually in the first 2 months and then remained unchanged. Compared with the invariant priming caused by repeated and continuous additions in the tropical soil, the priming induced by these addition patterns increased with time in the subtropical (Fig. 1e,f; Table S2). On an SOC-specific basis, priming induced by single addition was significantly higher in the tropical soil than in the subtropical in the first 3 months, but they were similar by the end of incubation. Finally, there were no significant differences in priming induced by repeated and continuous additions to these soils (Fig. 1e,f; Table S2).




Fig. 1 Incubation results for tropical (left) and subtropical (right) soils. Panels (a) and (b) show weekly total CO₂ fluxes in each treatment and soil. Panels (c) and (d) show the cumulative fraction of added glucose released as CO₂. Panels (e) and (f) show the cumulative net priming of SOC. Panels (g) and (h) show the net C balance between SOC priming and retention of added glucose. Color lines represent different treatments, i.e. black lines with empty circles in (a,b) indicate water-only, while red lines with triangles, blue lines with crossed squares, and pink lines with filled diamonds in (a-h) indicate single, repeated and continuous glucose additions to soils respectively. (c,d) indicate cumulative fraction of glucose released as CO₂, (e,f) indicate cumulative fluxes of SOC priming, and (g,h) indicate cumulative (positive) net effects of glucose additions on net soil C balances. Blue arrows in (e,f) indicate times of glucose additions to repeated treatments.

Net C balance of primed SOC and net retention of added glucose

The C retained from the glucose (not released as CO₂) exceeded the amount of primed SOC throughout the 170-day incubation period. Thus, net-C balances in both soils and all addition treatments were positive (Fig. 1g,h; Table S3; Tukey's HSD test, $P < 0.05$). Single additions to both soils initially caused significantly higher net-C increases than repeated and continuous additions. Positive net-C increase from single additions to tropical soil was constant, but it decreased through time in the subtropical soil (Fig. 1g,h; Table S3). At the end of the incubations, net-C increase was higher in continuous than single in the tropical soil (Fig. 1g). In the subtropical soil, net-C increases caused by single addition were significantly lower than in the other two addition patterns (Fig. 1h; Table S3). Net-C increases caused by continuous addition in the tropical soil were significantly higher than those caused by all three treatments in the subtropical soil (Fig. 1g,h; Table S3).

Responses of total CO₂ flux and primed C to added glucose

The first glucose additions to each soil varied in amount, but the amount was the same on an SOC-specific basis for the same treatment in both soils. Therefore, the correlation between initial CO₂ effluxes and glucose additions reflected microbial utilization for added glucose. The tropical-soil response was much larger (slopes of tropical vs. subtropical: 0.554 vs. 0.277; $P < 0.0001$; Fig. 2). Both soils responded linearly to glucose inputs ($R^2 = 0.99$; Fig. 2), indicating that microbial responses were not saturated even at the highest addition levels. The initial primed C of the tropical soil was probably higher (slopes of tropical vs. subtropical: 0.387 vs. 0.139; $P < 0.05$; Fig. 2). Since continuous and repeated glucose addition levels were much lower and closer to each other, the relatively close correlations presented here largely relied on the highest glucose additions. More intermediate values would make the correlation observed more convincing.

Microbial biomass, dissolved organic C and incorporation of glucose-C into microbial biomass

At the end of incubations, a significant decrease in DOC was observed only in repeated and continuous additions to the tropical soil, while there was no significant difference between single addition and the control treatment (Tukey's HSD test, $P < 0.05$, Fig. 3a). There were no treatment differences in the subtropical soil. Control, single, and repeated additions had higher MBC in subtropical than tropical soils (Fig. 3b). Continuous addition had higher MBC than other treatments in tropical soil, but there were no treatment differences in the subtropical soil (Tukey's HSD, $P < 0.05$, Fig. 3b).




Fig. 2 Responses of total CO₂ flux (top) and primed C (bottom) to the amount of added glucose in tropical and subtropical soils. Red long dash lines and empty circles indicate the tropical soil while blue solid lines and empty triangles represent the subtropical soil. There were 4 replicates for each treatment.




Fig. 3 Concentrations of dissolved organic C and microbial biomass in subtropical soil and tropical soils at the end of a 170-day incubation period. Bars show standard errors of the means ($n = 4$). Different letters above columns indicate significant difference among treatments for each soil at $P < 0.05$ level. The asterisk above columns indicates significant difference in microbial biomass C between subtropical and tropical soils at $P < 0.05$ level.

In both soils, the amount of primed SOC released as CO₂ was negatively correlated with glucose incorporation into microbial biomass ($R^2 = 0.63$, $P < 0.001$, Fig. 4), indicating that higher priming was accompanied by faster microbial turnover. Even though this relationship was similar for both soils, tropical soil was more variable in microbial glucose incorporation (range from 0.63 to 1.85% for tropical and from 0.76 to 1.37% for subtropical, Fig. 4).

Discussion

Rhizodeposition and decomposing litter DOC fluxes compared with experimental glucose additions

The glucose additions in this incubation study correspond to approximately 200 g C m⁻² yr⁻¹ to the soil surface. Rhizodeposition could represent




Fig. 4 Correlations between cumulative SOC priming and glucose incorporated and remaining in microbial biomass after 170-day incubation of subtropical and tropical soils. Open circles indicate the subtropical, and filled circles indicate the tropical soils.

100 g C m⁻² yr⁻¹, DOC from litter decomposition 10 g C m⁻² yr⁻¹, and future CO₂ doubling could increase total DOC fluxes by 50 g C m⁻² yr⁻¹ (see Introduction and Methods sections). None of those estimates are directly applicable to the forests we studied, but they suggest that we are exploring an appropriate range of LOC additions (de Graaff *et al.*, 2010).

Dependence of priming effects on C-addition patterns

To the best of our knowledge, this is the first time that priming effects induced by different addition patterns (i.e. single, repeated, and continuous additions) with the same total amounts have been compared in soils. We demonstrate that addition patterns can stimulate different priming of SOM decomposition, which strongly rely on initial addition amount, soil type, and the timing since glucose was added (Fig. 1e,f). The results of the initial glucose additions to both soils clearly showed that the amount of C added is an important driver for priming (Fig. 2). However, throughout the experiment, single additions caused more priming than repeated and continuous additions in both soils, while continuous and repeated additions produced similar priming (Fig. 1e,f). These results indicate that single glucose additions cannot represent SOC priming resulting from repeated or continuous glucose inputs. Therefore, our hypothesis that continuous glucose additions would induce more SOC priming than single additions was not supported.

The differences among addition patterns could be ascribed to different microbial activities, because the activation of microorganisms from no-growth or starvation states to active states (Morita, 1990; Blagodatskaya & Kuzyakov, 2008) by easily available organics can be

the main reason for accelerated SOM mineralization (Kuzyakov *et al.*, 2000; Hamer & Marschner, 2005; Kuzyakov, 2010). Most soil microorganisms are energy limited and oscillate between dormant and active physiological states (Morita, 1990; Stenström *et al.*, 2001). A fraction of microorganisms are activated when LOC arrives in soils (Stenström *et al.*, 2001). Microbial growth and enzyme production are accompanied by increased microbial demand for nutrients. Subsequently, microorganisms scavenge N from SOM and cause priming (Kuzyakov *et al.*, 2002; Fontaine *et al.*, 2004a; Cheng, 2009; Dijkstra *et al.*, 2013; Paterson & Sim, 2013). This is apparent in the subtropical soil but only for single additions in the tropical soil.

Although previous studies have demonstrated that microbial biomass can be activated by trace amounts of available organics and produce positive priming (De Nobili *et al.*, 2001; Mondini *et al.*, 2006), small continuous glucose additions were not enough to activate microorganisms and trigger positive priming in this tropical soil, despite increased microbial biomass compared with the control (Fig. 3). Even repeated glucose additions caused little priming. In contrast, small glucose additions to the subtropical soil triggered positive priming after the first month, and priming increased similarly as with repeated additions. This may result from alleviating microbial C or energy limitation, thus leading to increased microbial activities and nutrient demand (Cheng & Kuzyakov, 2005). These results indicate that the occurrence of positive priming requires the amount of added LOC to reach some threshold, but this threshold differs between soils (Fig. 1e,f). Several differences could contribute to this. First, we suggest that different physiological states of microorganisms in tropical and subtropical soils could be responsible for their activation by inputs of organics because inherent physiological states are strongly related to history (Stenström *et al.*, 2001). Compared with subtropical, the tropical SOC could be highly processed by microorganisms. Low MBC to SOC ratios from the controlled treatment of the tropical soil reflect high fraction of recalcitrant OC (Moscatelli *et al.*, 2005). Therefore, microorganisms could require more energy to decompose tropical SOC. Second, the availability of substrates in these soils (Fig. 3, top), including pools of available C and nutrients, could affect the direction and magnitude of priming induced by addition patterns. Numerous studies have demonstrated that microbial growth and growth states strongly depend on the size and quality of soil-available resources (Stenström *et al.*, 2001; Blagodatskaya & Kuzyakov, 2008).

Soils with higher C : N ratios generally exhibit stronger priming than those with low C:N ratios due to lack of available N (Zhang *et al.*, 2013), but our results did

not support this pattern. In our study, both soils produced similar priming effects under corresponding treatments by the end of incubations (Fig. 1e,f), and that single additions induced stronger priming for the first 3 months in the tropical soil having a lower C:N ratio (Table 1). We suggest vegetation, climate, microbial communities and soil properties other than C:N could be more important for the occurrence of priming. Therefore, our hypothesis that greater SOC priming would occur in the tropical soil than in the subtropical soil was only partly supported by our results.

Priming induced by a single glucose addition reached a maximum more rapidly in the tropical soils (Fig. 1e), suggesting more rapid microbial response to added glucose in the tropical soil (Paterson & Sim, 2013). A previous study also suggested that tropical soil C is very sensitive to changes in C inputs (Leff *et al.*, 2012). However, the underlying mechanisms should be investigated for a better understanding of SOC decomposition and turnover in forest soils.

Net soil C balance between SOC priming and net glucose retention

We observed that priming increased decomposition of SOC in both forest soils (Fig. 1e,f). However, the effects of LOC inputs on SOC dynamics should be evaluated in the context of net C balance. In the present study, we excluded CO₂ fluxes from the water-only treatments in calculating net C balances. These balances were always positive and initially (shortly after addition) highest with large glucose additions (Fig. 1g,h). High net-C accumulations in both soils with continuous glucose additions suggest that LOC inputs such as root exudation may increase SOC as observed by free-air CO₂ enrichments (Hoosbeek *et al.*, 2004; Liu *et al.*, 2009), even with soil C cycling accelerated by rhizosphere priming (Kuzyakov *et al.*, 2007; Cheng, 2009; Zhu & Cheng, 2011; Cheng *et al.*, 2013). Therefore, our hypothesis that the net C balances between primed C and the gain from added glucose could be negative in both forest soils was not supported.

To explore relationships between the amount of added glucose and the net SOC balance between primed C and the gain from added glucose, we compared our results with those of 13 previous studies (Fig. 5). Including the retention of added LOC, most of the studies showed net C increases in soil despite SOC priming (Fig. 5, top) and in only a very few studies priming was accompanied by a net decrease in SOC (Fig. 5, bottom). Some studies concluded that priming leads to SOC losses, but net SOC balance was usually positive when the remaining LOC was accounted for. Net soil C increases have been demonstrated in




Fig. 5 Net soil C balance under priming by additions of glucose and other labile organic carbon (LOC) from this and previous studies. Open circles represent our incubations of tropical and subtropical forest soils, filled circles represent data presented by Fontaine *et al.*, 2004a,b; Hamer & Marschner, 2005; Ohm *et al.*, 2007;. The X-symbols represent our calculations of soil-C balance from Wu *et al.*, 1993; Bell *et al.*, 2003; Brant *et al.*, 2006; Blagodatskaya *et al.*, 2007; Fontaine *et al.*, 2007; Hoyle *et al.*, 2008; Nottingham *et al.*, 2009; Guenet *et al.*, 2010; Garcia-Pausas & Paterson, 2011, based on our calculations from data presented by those authors. Positive net C balances are on the top panel and negative on the bottom.

other studies, e.g., free-air CO₂ enrichment, in which ¹³C-depleted CO₂ was added to small forest stands, and net soil C balances were examined. Most of these studies (Hoosbeek *et al.*, 2004; Liu *et al.*, 2009) found net SOC increases, while some observed a neutral balance (Jastrow *et al.*, 2005; van Kessel *et al.*, 2006) or net SOC decreases (Hoosbeek & Scarascia-Mugnozza, 2009). Similarly, Gattinger *et al.* (2012) reviewed studies on reduced-tillage agriculture and found that SOC increased, even though increased plant litter may have primed SOC losses.

Thus, several lines of evidence support our conclusion that net SOC often increases in response to C inputs, but particular local conditions may also affect C balances. Despite accelerated turnover by priming, increased C input contributes to C sequestration in the soil (Phillips *et al.*, 2011; Cheng *et al.*, 2013). Priming studies should consider net C-balances between primed C and the gain from added LOC and accelerated mineralization of native SOM is only a part of the C budget. It has also been suggested that priming can act on recalcitrant SOC (Fontaine *et al.*, 2007; Blagodatskaya *et al.*, 2011). If retained C from added LOC is less recalcitrant than primed C, then long-term effects on C sequestration remain uncertain. Therefore, further studies should

investigate the fate of added organics and sources of the primed C.

In summary, we found that single additions of glucose overestimate priming effects in these subtropical and tropical soils and thus do not adequately reflect natural ecosystem C inputs. Continuous C inputs resulted in a greater C increase in the soil than single additions, further emphasizing the importance of LOC addition patterns. Those patterns aside, responses of these very different soils to glucose additions were remarkably similar in terms of glucose release, priming, and net-C balance. From this experiment and previous studies, we present evidence that LOC additions may generally increase net C balances in soils. Accurate evaluation of the direction and intensity of priming must consider the observation period, dynamics of C utilization, and induced CO₂ losses, as well as the net C balance and other soil properties. Understanding and modeling effects of global change such as warming and elevated CO₂ on soil-C cycles need to be linked to plant production of LOC and its effects on microbial activity (Cheng *et al.*, 2013).

Acknowledgments

We thank Dr. Lu Qiao for her help in the field. We also thank Dr. Richard Corlett and Dr. Jonathan Adams for their helpful comments. This study was supported by the National Natural Science Foundation of China (41071209 and 30970535), the Knowledge Innovation Foundation Program for Outstanding Young Scholar of the Chinese Academy of Sciences (KZCX2-YW-QN302), and Strategic Pilot Program of Chinese Academy of Sciences (XDA05050203-02) and a Chinese Academy of Sciences Visiting Professorship to Yakov Kuzyakov. We also thank the anonymous reviewers for their substantial and constructive comments. We also thank Ailao-shan Station for Subtropical Forest Ecosystem Studies and Xishuangbanna Station for Tropical Forest Ecosystem Studies for providing background data.

References

- Allison SD, Wallenstein MD, Bradford MA (2010) Soil-carbon response to warming dependent on microbial physiology. *Nature Geoscience*, **3**, 336–340.
- Bell JM, Smith JL, Bailey VL, Bolton H (2003) Priming effect and C storage in semi-arid no-till spring crop rotations. *Biology and Fertility of Soils*, **37**, 237–244.
- Blagodatskaya EV, Kuzyakov Y (2008) Mechanisms of real and apparent priming effects and their dependence on soil microbial biomass and community structure: critical review. *Biology and Fertility of Soils*, **45**, 115–131.
- Blagodatskaya EV, Blagodatsky SA, Anderson TH, Kuzyakov Y (2007) Priming effects in Chernozem induced by glucose and N in relation to microbial growth strategies. *Applied Soil Ecology*, **37**, 95–105.
- Blagodatskaya E, Yuyukina T, Blagodatsky S, Kuzyakov Y (2011) Three sources partitioning of microbial biomass and CO₂ efflux from soil to evaluate mechanisms of priming effects. *Soil Biology & Biochemistry*, **43**, 778–786.
- Brant JB, Sulzman EW, Myrold DD (2006) Microbial community utilization of added carbon substrates in response to long-term carbon input manipulation. *Soil Biology & Biochemistry*, **38**, 2219–2232.
- Buitenwerf R, Bond WJ, Stevens N, Trollope SW (2012) Increased tree densities in South African savannas: >50 years of data suggests CO₂ as a driver. *Global Change Biology*, **8**, 675–684.

- Cernusak LA, Winter K, Dalling JW *et al.* (2013) Tropical forest responses to increasing atmospheric CO₂: current knowledge and opportunities for future research. *Functional Plant Biology*, **40**, 531–551.
- Chapin FS, Matson PA, Vitousek PM (2012) *Principles of Terrestrial Ecosystem Ecology*, 2nd edn. Springer, New York.
- Cheng W (1996) Measurement of rhizosphere respiration and organic matter decomposition using natural ¹³C. *Plant and Soil*, **183**, 263–268.
- Cheng WX (2009) Rhizosphere priming effect: Its functional relationships with microbial turnover, evapotranspiration, and C-N budgets. *Soil Biology & Biochemistry*, **41**, 1795–1801.
- Cheng WX, Kuzyakov Y (2005) Root effects on soil organic matter decomposition. In: *Roots and Soil Management: Interactions Between Roots and the Soil*. Agronomy Monograph no. 48. (eds Zobel RW, Wright SF), pp. 119–143. American Society of Agronomy, Crop Science Society of America, Soil Science Society of America. Madison, WI, USA.
- Cheng WX, Parton WJ, Gonzalez-Meler MA *et al.* (2013) Synthesis and modeling perspectives of rhizosphere priming. *New Phytologist*. doi: 10.1111/nph.12440.
- Chigineva NI, Aleksandrova AV, Tiunov AV (2009) The addition of labile carbon alters litter fungal communities and decreases litter decomposition rates. *Applied Soil Ecology*, **42**, 264–270.
- Cox PM, Pearson D, Booth BB, Friedlingstein P, Huntingford C, Jones CD, Luke CM (2013) Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability. *Nature*, **494**, 341–345.
- Davidson EA, Janssens IA (2006) Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature*, **440**, 165–173.
- De Nobili M, Contin M, Mondini C, Brookes PC (2001) Soil microbial biomass is triggered into activity by trace amounts of substrate. *Soil Biology & Biochemistry*, **33**, 1163–1170.
- Dieleman WJ, Luyssaert S, Rey A *et al.* (2010) Soil [N] modulates soil C cycling in CO₂-fumigated tree stands: a meta-analysis. *Plant, Cell and Environment*, **33**, 2001–2011.
- Dijkstra FA, Carrillo Y, Pendall E, Morgan JA (2013) Rhizosphere priming: a nutrient perspective. *Frontiers in Microbiology*, **4**, 1–8.
- Don A, Kalbitz K (2005) Amounts and degradability of dissolved organic carbon from foliar litter at different decomposition stages. *Soil Biology and Biochemistry*, **37**, 2171–2179.
- Drake JE, Gallet-Budynek A, Hofmocker KS *et al.* (2011) Increases in the flux of carbon belowground stimulate nitrogen uptake and sustain the long-term enhancement of forest productivity under elevated CO₂. *Ecology Letters*, **14**, 349–357.
- Drigo D, Kowalchuk GA, van Veen JA (2008) Climate change goes underground: effects of elevated atmospheric CO₂ on microbial community structure and activities in the rhizosphere. *Biology and Fertility of Soils*, **44**, 667–679.
- Drigo D, Kowalchuk GA, Knapp BA, Pijl AS, Boschker HTS, van Veen JA (2013) Impacts of 3 years of elevated atmospheric CO₂ on rhizosphere carbon flow and microbial community dynamics. *Global Change Biology*, **19**, 621–636.
- Falster DS, Wharton DJ, Wright IJ (2006). *SMATR: Standardised Major Axis Tests and Routines*. Available at: <http://www.bio.mq.edu.au/ecology/SMATR/> (accessed 13 November 2006).
- Finzi AC, Norby RJ, Calfapietra C, Callet-Budynek A, Gielen B *et al.* (2007) Increases in nitrogen uptake rather than nitrogen-use efficiency support higher rates of temperate forest productivity under elevated CO₂. *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 14014–14019.
- Fontaine S, Bardoux G, Abbadie L, Mariotti A (2004a) Carbon input to soil may decrease soil carbon content. *Ecology Letters*, **7**, 314–320.
- Fontaine S, Bardoux G, Benet D, Verdier B, Mariotti A, Abbadie L (2004b) Mechanisms of the priming effect in a savannah soil amended with cellulose. *Soil Science Society of American Journal*, **68**, 125–131.
- Fontaine S, Barot S, Barre P, Bdioui N, Mary B, Rumpel C (2007) Stability of organic carbon in deep soil layers controlled by fresh carbon supply. *Nature*, **450**, 277–281.
- Garcia-Pausas J, Paterson E (2011) Microbial community abundance and structure are determinants of soil organic matter mineralisation in the presence of labile carbon. *Soil Biology & Biochemistry*, **43**, 1705–1713.
- Gattinger A, Muller A, Haeni M, Skinner C, Fliessbach A *et al.* (2012) Enhanced top soil carbon stocks under organic farming. *Proceedings of the National Academy of Sciences of the United States of America*, **109**, 18226–18231.
- German DP, Chacon SS, Allison SD (2011) Substrate concentration and enzyme allocation can affect rates of microbial decomposition. *Ecology*, **92**, 1471–1480.
- de Graaff M-A, Classen AT, Castro HF, Schadt CW (2010) Labile soil carbon inputs mediate the soil microbial community composition and plant residue decomposition rates. *New Phytologist*, **188**, 1055–1064.
- Guenet B, Neill C, Bardoux G, Abbadie L (2010) Is there a linear relationship between priming effect intensity and the amount of organic matter input? *Applied Soil Ecology*, **46**, 432–442.
- Hamer U, Marschner B (2005) Priming effects in soils after combined and repeated substrate additions. *Geoderma*, **128**, 38–51.
- van Hees PAW, Jones DL, Finlay R, Godbold DL, Lundström US (2005) The carbon we do not see – the impact of low molecular weight compounds on carbon dynamics and respiration in forest soils: a review. *Soil Biology & Biochemistry*, **37**, 1–13.
- Heimann M, Reichstein M (2008) Terrestrial ecosystem carbon dynamics and climate feedbacks. *Nature*, **451**, 289–292.
- Hickler TS, Smith B, Prentice IC, Mjöfors K, Miller P, Arnett A, Sykes MT (2008) CO₂ fertilization in temperate FACE experiments not representative of boreal and tropical forests. *Global Change Biology*, **14**, 1531–1542.
- Hoosbeek MR, Scarascia-Mugnozza GE (2009) Increased litter build up and soil organic matter stabilization in a poplar plantation after 6 years of atmospheric CO₂ enrichment (FACE): final results of POP-EuroFACE compared to other forest FACE experiments. *Ecosystems*, **12**, 220–239.
- Hoosbeek MR, Lukac M, van Dam D, Godbold DL, Velthorst EJ *et al.* (2004) More new carbon in the mineral soil of a poplar plantation under free air carbon enrichment (POPFACE): Cause of increased priming effect? *Global Biogeochemical Cycles*, **18**, GB1040, doi: 10.1029/2003GB002127.
- Hoyle FC, Murphy DV, Brookes PC (2008) Microbial response to the addition of glucose in low-fertility soils. *Biology and Fertility of Soils*, **44**, 571–579.
- Hungate BA, Van Groenigen K-J, Six J *et al.* (2009) Assessing the effect of elevated carbon dioxide on soil carbon: a comparison of four meta-analyses. *Global Change Biology*, **15**, 2020–2034.
- Jackson RB, Cook CW, Pippin JS, Palmer SM (2009) Increased belowground biomass and soil CO₂ fluxes after a decade of carbon dioxide enrichment in a warm-temperate forest. *Ecology*, **90**, 3352–3366.
- Jastrow JD, Miller RM, Matamala R, Norby RJ, Boutton TW *et al.* (2005) Elevated atmospheric carbon dioxide increases soil carbon. *Global Change Biology*, **11**, 2057–2064.
- Jones DL, Darrah PR (1993) Re-sorption of organic compounds by roots of *Zea mays* L. and its consequences in the rhizosphere. II. Experimental and model evidence for simultaneous exudation and re-sorption of soluble C compounds. *Plant and Soil*, **153**, 47–59.
- van Kessel C, Horwarth WR, Hartwig U, Harris D, Lüscher A (2006) Net soil carbon input under ambient and elevated CO₂ concentrations: isotopic evidence after 4 years. *Global Change Biology*, **6**, 435–444.
- Kirschbaum MUF (2004) Soil respiration under prolonged soil warming: are rate reductions caused by acclimation or substrate loss? *Global Change Biology*, **10**, 1–8.
- Krull ES, Skjemstad JO (2003) $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ profiles in ¹⁴C-dated Oxisol and Vertisols as a function of soil chemistry and mineralogy. *Geoderma*, **112**, 1–29.
- Ku H (1966) Notes on the Use of Propagation of Error Formulas, Journal Research of National Bureau of Standards-C. *Engineering and Instrumentation*, **70C**, 263–273.
- Kuzyakov Y (2010) Priming effects: interactions between living and dead organic matter. *Soil Biology & Biochemistry*, **42**, 1363–1371.
- Kuzyakov Y, Friedel JK, Stahr K (2000) Review of mechanisms and quantification of priming effects. *Soil Biology & Biochemistry*, **32**, 1485–1498.
- Kuzyakov Y, Siniakina SV, Rühlmann J, Domanski G, Stahr K (2002) Effect of nitrogen fertilisation on below-ground carbon allocation in lettuce. *Journal of the Science of Food and Agriculture*, **82**, 1432–1441.
- Kuzyakov Y, Hill PW, Jones DL (2007) Root exudate components change litter decomposition in a simulated rhizosphere depending on temperature. *Plant and Soil*, **290**, 293–305.
- Leff JW, Wieder WR, Taylor PG, Townsend AR, Nemergut DR, Grandy AS, Cleveland CC (2012) Experimental litterfall manipulation drives large and rapid changes in soil carbon cycling in a wet tropical forest. *Global Change Biology*, **18**, 2969–2979.
- Lewis SL, Lloyd J, Sitch S, Mitchard TA, Laurance WF (2009) Changing ecology of tropical forests: evidence and drivers. *Annual Review of Ecology, Evolution, and Systematics*, **40**, 529–549.
- Li H (1983) Ecological characteristics of vegetation in Xujiaaba region of Ailao Mts. In: *Research of Forest Ecosystem on Ailao Mountains Yunnan* (ed. Li H), pp. 63–73. Yunnan Science and Technology Press, Kunming, China.
- Liski J, Nissinen A, Erhard M, Taskinen O (2003) Climatic effects on litter decomposition from arctic tundra to tropical rainforest. *Global Change Biology*, **9**, 575–584.
- Liu W, Fox JED, Xu ZF (2000) Leaf litter decomposition of canopy trees, bamboo and moss in a montane moist evergreen broad-leaved forest in Ailao Mountain, Yunnan, south-west China. *Ecological Research*, **15**, 435–447.

- Liu LL, King JS, Booker FL, Giardina CP, Allen HL, Hu SJ (2009) Enhanced litter input rather than changes in litter chemistry drive soil carbon and nitrogen cycles under elevated CO₂: a microcosm study. *Global Change Biology*, **15**, 441–453.
- Meharg AA, Kilham K (1991) A novel method of quantifying root exudation in the presence of soil microflora. *Plant and Soil*, **113**, 111–116.
- Mondini C, Cayuela ML, Sanchez-Monedero MA, Roig A, Brookes PC (2006) Soil microbial biomass activation by trace amounts of readily available substrate. *Biology and Fertility of Soils*, **42**, 542–549.
- Morita RY (1990) The starvation-survival state of microorganisms in nature and its relationship to the bioavailable energy. *Cellular and Molecular Life Sciences*, **46**, 813–817.
- Moscattelli M, Lagomarsino A, Marinari S, De Angelis P, Grego S (2005) Soil microbial indices as bioindicators of environmental changes in a poplar plantation. *Ecological Indicators*, **5**, 171–179.
- Nottingham AT, Griffiths H, Chamberlain PM, Stott AW, Tanner EVJ (2009) Soil priming by sugar and leaf-litter substrates: A link to microbial groups. *Applied Soil Ecology*, **42**, 183–190.
- Ohm H, Hamer U, Marschner B (2007) Priming effects in soil size fractions of a podzol Bs horizon after addition of fructose and alanine. *Journal of Plant Nutrition and Soil Science*, **170**, 551–559.
- Pallant J (2007) *SPSS Survival Manual: a Step by Step Guide to Data Analysis Using SPSS for Windows*, Open University Press, McGraw-Hill, Maidenhead, Berkshire.
- Paterson E, Sim A (2013) Soil-specific response functions of organic matter mineralization to the availability of labile carbon. *Global Change Biology*, **19**, 1562–1571.
- Pendall E, Bridgman S, Hanson PJ *et al.* (2004) Below-ground process responses to elevated CO₂ and temperature: a discussion of observations, measurement methods, and models. *New Phytologist*, **162**, 311–322.
- Phillips DL, Gregg JW (2001) Uncertainty in source partitioning using stable isotopes. *Oecologia*, **127**, 171–179.
- Phillips DL, Newsome SD, Gregg JW (2005) Combining sources in stable isotope mixing models: alternative methods. *Oecologia*, **144**, 520–527.
- Phillips RP, Ehlitz Y, Bier R, Bernhardt ES (2008) New approach for capturing soluble root exudates in forest soils. *Functional Ecology*, **22**, 990–999.
- Phillips RP, Bernhardt ES, Schlesinger WH (2009) Elevated CO₂ increases root exudation from loblolly pine (*Pinus taeda*) seedlings as an N-mediated response. *Tree Physiology*, **29**, 1513–1523.
- Phillips RP, Finzi AC, Bernhardt ES (2011) Enhanced root exudation induces microbial feedbacks to N cycling in a pine forest under long-term CO₂ fumigation. *Ecology Letters*, **14**, 187–194.
- R Development Core Team (2011) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Raich JW, Potter CS, Bhagawati D (2002) Interannual variability in global soil respiration, 1980–1994. *Global Change Biology*, **8**, 800–812.
- Raich JM, Russell AE, Kitayama K, Parton W, Vitousek PM (2006) Temperature influences carbon accumulation in moist tropical forests. *Ecology*, **87**, 76–87.
- Reich PB (2011) Taking stock of forest carbon. *Nature Climate Change*, **1**, 346–347.
- Rumpel C, Kögel-Knabner I (2011) Deep soil organic matter—a key but poorly understood component of terrestrial C cycle. *Plant and Soil*, **338**, 143–158.
- Sayer EJ, Heard MS, Grant HK, Marthews TR, Tanner EVJ (2011) Soil carbon release enhanced by increased tropical forest litterfall. *Nature Climate Change*, **1**, 304–307.
- Schaefer DA, Feng WT, Zou XM (2009) Plant carbon inputs and environmental factors strongly affect soil respiration in a subtropical forest of southwestern China. *Soil Biology & Biochemistry*, **41**, 1000–1007.
- Stenström J, Svensson K, Johansson M (2001) Reversible transition between active and dormant microbial states in soil. *FEMS Microbiology Ecology*, **36**, 93–104.
- Tan Z, Zhang YP, Yu GR, Sha L, Tang J, Deng X, Song Q (2010) Carbon balance of a primary tropical seasonal rain forest. *Journal of Geophysical Research*, **115**, D00H26, doi: 10.1029/2009JD012913.
- Tan ZH, Zhang YP, Schaefer D, Yu GR, Liang N, Song QH (2011) An old-growth subtropical Asian evergreen forest as a large carbon sink. *Atmospheric Environment*, **45**, 1548–1554.
- Tang J-W, Cao M, Zhang J-H, Li M-H (2010) Litterfall production, decomposition and nutrient use efficiency varies with tropical forest types in Xishuangbanna, SW China: a 10-year study. *Plant and Soil*, **335**, 271–288.
- Tarnocai C, Canadell JG, Schuur EAG, Kuhry P, Mazhitova G, Zimov S (2009). Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochemical Cycles*, **23**, GB2023, doi: 10.1029/2008GB003327.
- Trofymow JA, Coleman DC, Cambardella C (1987) Rates of rhizodeposition and ammonium depletion in the rizosphere of axenic oat roots. *Plant and Soil*, **97**, 333–344.
- Vance ED, Brookes PC, Jenkinson DS (1987) An extraction method for measuring soil microbial biomass C. *Soil Biology & Biochemistry*, **19**, 703–707.
- Warton DI, Wright IJ, Falster DS, Westoby M (2006) A review of bivariate line-fitting methods for allometry. *Biological Reviews*, **81**, 259–291.
- Wood TE, Cavaleri MA, Reed SC (2012) Tropical forest carbon balance in a warmer world: a critical review spanning microbial- to ecosystem-scale processes. *Biological Reviews*, **87**, 912–927.
- World Reference Base for Soil Resources (2006) *World Soil Resources Reports 103*. FAO, Rome.
- Wu ZY (1980) *Vegetation of China*. Science Press, Beijing.
- Wu J, Brookes PC, Jenkinson DS (1993) Formation and destruction of microbial biomass during the decomposition of glucose and ryegrass in soil. *Soil Biology & Biochemistry*, **25**, 1435–1441.
- Wu Z, Dijkstra P, Koch GW, Peñuelas J, Hungate BA (2011) Responses of terrestrial ecosystems to temperature and precipitation change: a meta-analysis of experimental manipulation. *Global Change Biology*, **17**, 927–942.
- Würth MKR, Winter K, Körner C (1998) *In situ* responses to elevated CO₂ in tropical forest understorey plants. *Functional Ecology*, **12**, 886–895.
- Zhang K (1983) The characteristics of mountain climate in the north of Ailao Mts. *Research of Forest Ecosystem on Ailao Mountains, Yunnan*, (ed. Wu ZY), pp. 20–29. Yunnan Science and Technology Press, Kunming, China.
- Zhang WD, Wang XF, Wang SL (2013) Addition of external organic carbon and native soil organic carbon decomposition: a meta-analysis. *PLoS ONE*, **8**, e54779. doi: 10.1371/journal.pone.0054779.
- Zhu B, Cheng WX (2011) Rhizosphere priming effect increases the temperature sensitivity of soil organic matter decomposition. *Global Change Biology*, **17**, 2172–2183.
- Zibilske LM (1994) Carbon mineralization. In: *Methods of Soil Analysis, Part 2, Microbiological and Biochemical Properties* (eds Weaver RW, Angle S, Bottomley P *et al.*), pp. 835–863. Wisconsin, USA, Soil Science Society of America.
- Ziska LH, Hogan KP, Smith AP, Drake BG (1991) Growth and photosynthetic response of nine tropical species with long-term exposure to elevated carbon dioxide. *Oecologia*, **86**, 383–389.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Results of Tukey's HSD test on cumulative glucose CO₂ (as % of added glucose), cumulative priming effect and net carbon balance among different treatments and those between two forest soils, which were shown in Fig. 1.

Table S1. Results of Tukey's HSD test on cumulative glucose CO₂ (as% of added glucose) among different treatments and those between two forest soils.

Table S2. Results of Tukey's HSD test on cumulative priming effect among different treatments and those between two forest soils.

Table S3. Results of Tukey's HSD test on net carbon balance among different treatments and those between two forest soils.