# New Indole Alkaloids from the North Sea Bacterium *Vibrio* parahaemolyticus Bio249<sup>1</sup>

Ravikanth Veluri<sup>†</sup>, Imelda Oka<sup>†</sup>, Irene Wagner-Döbler<sup>‡</sup>, and Hartmut Laatsch<sup>\*,†</sup>

Department of Organic Chemistry, University of Göttingen, Tammannstrasse 2, D-37077 Göttingen, Germany and Department of Microbiology at the National Research Institute for Biotechnology (GBF), Mascheroder Weg 1, D-38124 Braunschweig, Germany

(Received June \_\_\_\_, 2003)

<sup>\*</sup> To whom correspondence should be addressed. Tel.: +(551) 393211; Fax: +(551) 399660. e-mail: hlaatsc@gwdg.de.

<sup>&</sup>lt;sup>†</sup> University of Göttingen

<sup>&</sup>lt;sup>‡</sup> GBF Braunschweig

<sup>\*</sup> To whom correspondence should be addressed. Tel.: +(551) 393211; Fax: +(551) 399660. e-mail: hlaatsc@gwdg.de.

Three new bis- and tris-indole derivatives, 3,3-bis(3-indolyl)-butane-2-one (**1a**), 1,1,1-tris(3-indolyl)methane (**2a**) and 1,1,3-tris(3-indolyl)butane (**3**) were isolated from a North Sea bacterium which was closely related to *Vibrio parahaemolyticus* (97 % homology). Additionally, arundine (**1b**), a known indole from a plant source along with many other known compounds were isolated from the ethyl acetate extract of the culture. Their structures were established on the basis of various spectral data. All the new compounds were inactive against a range of bacteria and fungi.

More than 1000 alkaloids with the indole skeleton have been reported from microorganisms.<sup>2,3</sup> One third of these compounds are peptides with masses beyond 600 Dalton where the indole is tryptophan-derived. The structural variety of the other part is higher, and their biological activity seems to cover a broader range, including antimicrobial, antiviral, cytotoxic, insecticidal, antithrombotic or enzyme inhibitory activity to list only some of them.<sup>3</sup> In contrast, most of the peptides have mainly antibiotic activity. In our search for new bioactive molecules from marine microorganisms,<sup>4,5</sup> we isolated two new indole alkaloids, namely 3,3-bis(3-indolyl)-butane-2-one (**1a**) and 1,1,3-tris(3-indolyl)butane (**3**). Additionally, arundine<sup>6</sup> (**1b**), 1,1,1-tris(3-indolyl)methane<sup>7</sup> (**2a**) and several other metabolites have been found which have previously been obtained by synthesis or were reported from other sources. We report here one of the cases where the differentiation between a biosynthetic and an abiotic origin of the metabolites is difficult.

## **Results and Discussion**

The producing strain Bio249 was isolated from a biofilm grown on a glass plate in the North Sea and taxonomically classified as closely related to *Vibrio parahaemolyticus* on the basis of the 16S rRNA (98 % homology). On a casein medium, the strain did not produce antibiotic activities, however several UV absorbing compounds were formed which were tentatively identified as indoles due to their color reaction with Ehrlich's reagent. The ethyl acetate extract of a bulk culture of the microorganism was subjected to column chromatography on silica gel and Sephadex LH-20, which afforded a high yield of unsubstituted indole along with indole-3-carboxylic acid,<sup>8</sup> indole-3-carbaldehyde,<sup>8</sup> arundine<sup>6</sup> (**1b**), vibrindole A<sup>9</sup> (**1c**), trisindoline<sup>10</sup> (**4a**), 2,2-di(3-indolyl)-3-indolone<sup>9</sup> (**5**), pharacine,<sup>11</sup> *p*-hydroxy-phenylethanol,<sup>12</sup> phenyl acetamide,<sup>13</sup> and uracil. The indoles **1c** and **5** had been isolated from another strain of *V. parahaemolyticus* before,<sup>9</sup> and **4a** was known as a metabolite from a *Vibrio* sp. isolated from a sponge.<sup>10</sup> This is the first reported occurrence, however, of the plant metabolite arundine (**1b**) and of the synthetic tris-indole **2a** in a microorganism. The structures of all metabolites were identified by comparison with the reported spectral data (<sup>1</sup>H, <sup>13</sup>C and MS) using AntiBase.<sup>3</sup>

Compound **1a** was obtained as a pale yellow solid which gave a colorless spot on TLC. Its EI mass spectrum gave a molecular ion at 302 [ $M^{+}$ ]. The HREIMS data (302.1419) suggested that the molecular formula is C<sub>20</sub>H<sub>18</sub>N<sub>2</sub>O, and the IR spectrum showed characteristic bands at 3400 (NH) and 1694 (C=O) cm<sup>-1</sup>. Its <sup>1</sup>H NMR spectrum showed two broadened *o*-coupled

doublets at  $\delta$  7.40 and 7.38 (both J = 7.8 Hz), and two triplets at 7.17 and 7.02 (each J = 7.2 Hz) indicated the presence of a 1,2-disubstituted benzene ring. Another proton in the aromatic region at  $\delta$  6.94 (d, J = 2.2 Hz) together with the positive Ehrlich reaction on TLC indicated a 3-substituted indole ring. Furthermore its <sup>1</sup>H NMR spectrum showed the presence of two methyl singlets at  $\delta$  2.08 and 2.16. All the aromatic signals related to the indole system were found to be doubled in their intensity with respect to the methyl signals. The <sup>13</sup>C NMR and APT data indicated the presence of only 12 carbon signals of which five were due to quaternary carbon atoms, five methine carbons and two methyl groups. The <sup>1</sup>H NMR spectrum, the MS fragmentation pattern and high resolution MS confirmed the presence of two indolyl moieties in a symmetrical orientation. From the foregoing spectral data the structure of the compound **1a** was established as 3,3-bis(3-indolyl)-butane-2-one, which was obtained earlier synthetically.<sup>14</sup>

Compound **2a** was isolated as a strongly UV absorbing solid with a high resolution mass (361.1579) indicating the molecular formula  $C_{25}H_{19}N_3$ . The doublets at  $\delta$  7.48, 7.36 and 6.78, and two triplets at  $\delta$  7.18 and 6.98 in the <sup>1</sup>H NMR spectrum revealed the presence of the same 3-substituted indolyl system as found in the previous compound **1a**. Further, its <sup>1</sup>H NMR spectrum displayed a singlet at  $\delta$  6.18. All the aromatic signals were found to be tripled in their intensity relative to the higher field signal at  $\delta$  6.18. The <sup>13</sup>C and APT experiments indicated a total of only 9 carbon signals which included 6 methine and 3 quaternary carbon atoms in the skeleton. Except for the methine signal at  $\delta$  32.7, all the remaining signals were found to be due to the 3-substituted indolyl system. The low number of carbon signals together with the high mass suggested the structure to be highly symmetrical. According to the EI mass fragmentation pattern with a signal at  $\delta$  117 typical of 3-substituted indoles and an M-118 ion at 243, the structure of the compound was found to be 1,1,1-tris(3-indolyl)methane (**2a**).

## ((insert formulae 1-5 here))

The trimeric indole **2a** has not been reported so far, but a closely related compound, turbomycin A (**2b**), has been published recently as a natural product: Gillespie et al.<sup>15</sup> isolated DNA fragments from soil samples and expressed the genes in *E. coli*. One of the clones produced the orange **2b** which had been synthesized previously,<sup>7</sup> and the related turbomycin B. Both turbomycins exhibited broad spectrum antibiotic activity against both gram positive and gram negative bacteria whereas the compound **2a** was found to be inactive against *Escherichia coli*, *Staphylococcus aureus*, *Bacillus subtilis*, *Streptomyces viridochromogenes*,

*Mucor miehei*, and *Candida albicans*. The didehydro derivative of **2a**, bis(3-indolyl)-3*H*-indolylidenmethane, was formed by *Saccharomyces cerevisiae* under the influence of light.<sup>16</sup>

Compound **3** gave a high resolution EI mass spectrum (403.2030) suggesting the molecular formula  $C_{28}H_{25}N_3$ . The <sup>1</sup>H NMR spectrum showed typical aromatic signals indicating again the presence of the indolyl skeleton as observed in the previous compounds. From the <sup>1</sup>H, <sup>1</sup>H COSY, <sup>13</sup>C and HMQC data it was derived that the compound contained three indolyl fragments substituted at position 3. Further, its <sup>1</sup>H NMR spectrum showed two methine proton signals at  $\delta$  4.50 (t, *J* = 7.8 Hz) and 3.00 (m), a methylene signal at  $\delta$  2.40 (m) and 2.60 (m) and a methyl signal at  $\delta$  1.38 (d, *J* = 7.0 Hz). From the coupling patterns the partial structures CH-CH<sub>3</sub> and CH-CH<sub>2</sub>-CH were obtained and finally the COSY correlations indicated the presence of a trisubstituted butane fragment. From the HMQC and HMBC experiments the three indolyl fragments must be attached to the positions 1,1,3 of the butane backbone. In the HMBC spectrum, the methine proton at  $\delta$  4.50 showed correlations with the carbon atom 2 of the butane chain and the atoms 2", 3", 3a", 2"', 3"' and 3a''' of two indole moieties. Further important correlations are reported in Figure 1. The foregoing spectral data in addition to the EI mass fragmentation revealed the structure of compound **3** as 1,1,3-tris(3-indolyl)butane.

### ((insert figure 1 here))

## Figure 1. Selected HMBC correlations of 1,1,3-tris(3-indolyl)butane (3)

Although there are many reports on the antibiotic activities even of simple indole derivatives, we could not find any antibiotic or antifungal properties of the indole derivatives described here. 3-Hydroxymethyl indole, bis(3-indolyl)methane (**1b**) and many related compounds show, however, beside other effects pronounced antiproliferative properties<sup>17</sup> by a G1 cell cycle arrest<sup>18</sup> and were therefore suggested for the chemoprevention and treatment of cancer,<sup>19</sup> for the treatment of premenstrual syndrome,<sup>20</sup> or as anti-inflammatory agents, to list some of their activities.<sup>21</sup> The new indoles from *V. parahaemolyticus* might have related properties and will be investigated in this respect.

As indole reacts smoothly with carbonyl compounds affording the corresponding bisarylated products, the bisindolylmethane (**1b**) can be obtained easily by reaction of indole with formaldehyde under slightly acidic conditions.<sup>22</sup> In a similar manner, the reaction of indole-3-carbaldehyde with indole, both isolated from *V. parahaemolyticus*, should yield **2a**, and also all other bis- and tris-indoles isolated here may be derived by Pictet-Spengler type reactions from indole and the respective carbonyl compound. This pointed to the possibility that at least some of the metabolites might have been formed artificially from indole in a reaction catalyzed by traces of acetic acid liberated by hydrolysis of the moist ethyl acetate during work-up. Natural products with chiral elements are mostly found as optically active isomers, not as racemic mixtures. The compound **3**, however, did not show any CD effect and was racemic indeed which supported the assumption of a non-enzymatic origin.

As we did not find a way to isolate the potential low-molecular carbonyl precursors (formaldehyde, acetaldehyde, crotonaldehyde, diacetyl, isatin) directly from the crude extract, we added 2-methylindole before work-up of the culture broth to induce eventually the formation of the homologous methyl indole derivatives, e.g. **4b**. Indeed, the labelling experiment afforded additional HPLC signals in the crude extract at slightly higher retention times which showed mass differences of  $\Delta m/z = 14$  and 28 with respect to the original indole derivatives. In the case of the trisindoline (**4a**), MSMS of the molecular ion at m/z 363 gave only a M – 117 fragment corresponding to the loss of one indole unit. In the labelling experiment, a satellite signal with a slightly higher retention time and the mass of 377 showed fragment ions at 245 and 259 corresponding to the loss of either indole or methylindole as expected for a monomethylated trisindoline thus confirming that at least a part of **4a** was postbiosynthetically formed.

In an additional experiment, the bacterial cell mass was scraped off directly from the agar plate (pH 8.8), extracted by trituration with ethyl acetate and immediately analyzed by HPLC/MS. Although the concentration of the bis- and tris-indoles found was lower with respect to the indole concentration in the previous batch fermentation, arundine (1b) and the indoles 2a, 4a and 5 could be detected unequivocally.

Formaldehyde, acetaldehyde, and crotonaldehyde are common microbial metabolites. Isatin was isolated from a marine *Alteromonas* sp.,<sup>23</sup> and diacetyl is one of the odor components of the fungus *Zostera marina*.<sup>24</sup> From bacteria, diacetyl has not been described so far, however, extracts of marine Streptomycetes possessed sometimes a very characteristic buttery smell pointing to diacetyl, isolate B7064 being a recent example in our group.<sup>25</sup> Although we were not able to prove the presence of this ketone by spectroscopic methods, the typical flavor was a strong hint that diacetyl is also a bacterial metabolite. It seems clear therefore that a part of the condensation products is already formed during the life cycle of *Vibrio parahaemolyticus* Bio249, however, work-up conditions are also strongly involved in the formation of the different indole oligomers.

### **Experimental Section**

**General Experimental Procedures**. NMR spectra were measured on Varian Unity 300 (300.145 MHz) and a Varian Inova 500 (499.876 MHz) spectrometer in CDCl<sub>3</sub> with TMS as internal standard. EIMS was recorded on a Finnigan MAT95 (70 eV) and perfluorokerosene was used as reference substance in HREIMS. HPLC/ESI-MS was recorded on a Finnigan LCQ with quarternary pump Rheos 4000 (Flux Instrument) and nucleosil column EC 125/2, 100-5, C 18. IR spectra were recorded on a Perkin-Elmer 1600 Series FTIR in KBr pellets. Flash chromatography was carried out on silica gel (30-60 mm, J. T. Baker). *R*<sub>f</sub> values were measured on Polygram SIL G/UV<sub>254</sub> (Macherey-Nagel & Co., Düren, Germany) with cyclohexane/ethyl acetate 7:3. Size exclusion chromatography was done on Sephadex LH-20 (Pharmacia). The strain *Vibrio parahaemolyticus* isolate Bio249 is deposited in the culture collection of the Department of Microbiology at the National Research Institute for Biotechnology (GBF) in Braunschweig, Germany.

**Casein medium**: Casein (10 g), yeast extract (5 g), NaCl (10 g) and glucose (5 g) were dissolved in 1 L of tap water and the medium was adjusted to pH 7.8 with 2 N NaOH and sterilized for 33 minutes at 121 °C. After sterilization, an end pH of 6.5 of the medium was obtained.

**Fermentation and Isolation**: A 1 cm<sup>2</sup> piece of agar from a 7-day old culture of *Vibrio parahaemolyticus* isolate Bio249 grown on M<sub>2</sub> medium was used to inoculate 250 mL of casein medium in 1 L Erlenmeyer flasks on a total of 25 L. These cultures were incubated on a rotary shaker (180 rpm) at 30 °C for 4 days. The culture broth was mixed with diatom earth (1 kg) and filtered through a pressure filter. The culture filtrate and the mycelium were separately extracted each with ethyl acetate. The mycelium was additionally extracted with acetone. The combined organic phases were evaporated to dryness to obtain 3.1 g of crude extract. Thus obtained extract was subjected to column chromatography on silica gel with a stepwise CH<sub>2</sub>Cl<sub>2</sub>/MeOH gradient (0.5 L of CH<sub>2</sub>Cl<sub>2</sub>, 0.5 L of CH<sub>2</sub>Cl<sub>2</sub>/2 % MeOH, 0.5 L of CH<sub>2</sub>Cl<sub>2</sub>/15 % MeOH, 0.5 L CH<sub>2</sub>Cl<sub>2</sub>/7.5 % MeOH, 0.5 L CH<sub>2</sub>Cl<sub>2</sub>/10 % MeOH, 0.5 L of CH<sub>2</sub>Cl<sub>2</sub>/15 % MeOH, 0.3 L of CH<sub>2</sub>Cl<sub>2</sub>/20 % MeOH and 0.3 L of CH<sub>2</sub>Cl<sub>2</sub>/50 % MeOH) to deliver eight fractions. Further purification of fraction 1 on Sephadex LH-20 (3 × 60 cm, MeOH) yielded 200 mg of indole ( $R_f = 0.75$ ), 55 mg of 1,1,3-tris(3-indolyl)butane (**3**,  $R_f =$  0.3) and 3 mg of pharacine ( $R_f = 0.45$ ). Fraction 2 yielded 6 mg of 2,2-di(3-indolyl)-3indolone (**5**,  $R_f = 0.55$ ), 4 mg of vibrindole (**1c**,  $R_f = 0.5$ ), 6 mg of tris(3-indolyl)methane (**2a**,  $R_f = 0.38$ ) and 5 mg of trisindoline (**6a**,  $R_f = 0.2$ ). Fraction 3 and 4 yielded 9 mg of 1,1-bis(3indolyl)-butane-2-one (**1a**,  $R_f = 0.35$ ) and 4 mg of 1,1-di(3-indolyl)methane (**1b**,  $R_f = 0.48$ ) respectively. Fraction 5 and 6 yielded 10 mg of indole-3-carboxylic acid, 12 mg of indole-3carbaldehyde, 22 mg of *p*-hydroxy phenylethanol, 13 mg of phenylacetamide and 40 mg of thymine.

**3,3-Bis(3-indolyl)-butan-2-one** (**1a**): Slightly yellow solid; UV (MeOH)  $\lambda_{max}$  (log  $\varepsilon$ ) 274 sh (2.91), 281 (2.94), 289 (2.89); IR (KBr)  $\nu_{max}$  3423, 1694, 1636, 1455, 1422, 1339, 1245, 1200, 1111, 745 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz):  $\delta$  8.15 (s br, 2 NH), 7.40 (2 d, *J* = 7.8 Hz; 2 H, H-4,4'), 7.38 (2 d, *J* = 7.8 Hz; 2 H, H-7,7'), 7.17 (td, *J* = 7.2, *J* = 1.0 Hz; 2 H, H-5,5'), 7.02 (td, *J* = 7.2, *J* = 1.0 Hz; 2 H, H-6,6'), 6.94 (d, *J* = 2.2 Hz; 2 H, H-2,2'), 2.16 (s, 3 H, COMe), 2.08 (s, 3 H, Me); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75.5 MHz)  $\delta$  209.6 (s, Ac), 136.8 (s, C-7,7'a), 126.1 (s, C-3,3'a), 123.0 (d, C-2,2'), 122.0 (d, C-6,6'), 121.4 (d, C-4,4'), 119.5 (d, C-5,5'), 118.8 (s, C-3,3'), 111.3 (d, C-7,7'), 52.7 (s, C-1), 26.6 (q, Ac-Me), 24.8 (q, C-2); EIMS (70 eV) *m/z* 302, 259, 142, 115, 77; (-)-ESIMS *m/z* 301.8 [M-H<sup>+</sup>], 325 [M+Na<sup>+</sup>]; HREIMS *m/z* 302.1421 (calcd. 302.14191 for C<sub>20</sub>H<sub>18</sub>N<sub>2</sub>O).

**1,1,1-Tris(3-indolyl)methane** (**2a**): Yellowish solid; UV (MeOH)  $\lambda_{max}$  (log ε) 248 sh (3.56), 283 (3.32), 388 (3.65); IR (KBr)  $\nu_{max}$  3425, 2924, 2855, 1620, 1482, 1412, 1216, 1127, 747 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.90 (s br, 3 NH), 7.48 (d, *J* = 8.0 Hz, 3 H-4,4',4"), 7.36 (d, *J* = 8.0 Hz, 3 H, H-7,7',7"), 7.18 (td, *J* = 7.6, *J*' = 1.0 Hz, H-6,6',6"), 6.98 (td, *J* = 7.6, *J*' = 0.8 Hz, H-5,5',5"), 6.76 (d, *J* = 2.2 Hz, 3 H, H-2,2',2"), 6.18 (s, 1 H, H-1); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75.5 MHz) δ 138.5 (s, C-7'a), 128.6 (s, C-3'a), 124.5 (d, C-2'), 121.9 (d, C-6'), 120.6 (d, C-4'), 120.2 (s, C-3'), 119.2 (d, C-5'), 112.1 (d, C-7'), 32.7 (d, C-1); all dashed atoms are of threefold intensities; EIMS (70 eV) *m/z* 361 (100), 277 (10), 243 (50), 216 (15), 180 (10), 117 (18); HREIMS *m/z* 361.1577 (calcd. 361.15789 for C<sub>25</sub>H<sub>19</sub>N<sub>3</sub>).

**1,1,3-Tris(3-indolyl)butane** (**3**): Yellowish solid; UV (MeOH)  $\lambda_{max}$  (log ε) 276 sh (4.00), 282 (4.03), 290 (3.99); IR (KBr)  $\nu_{max}$  3411, 3051, 2924, 1620, 1456, 1419, 1337, 1219, 1094, 1011, 742, 582, 423 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.60 – 7.43 (m, 4 H, 3 NH, H-4'),

7.43 (d, J = 7.8 Hz, 1 H, H-4"), 7.41 (d, J = 7.8 Hz, 1 H, H-4"), 7.25 – 7.05 (m, 6 H-ar), 7.04 – 6.90 (m, 3 H, Ar-H), 6.76 (d, J = 2.0 Hz, 1 H, H-2"), 6.70 (d, J = 2.0 Hz, 1 H, H-2'), 6.65 (d, J = 2.0 Hz, 1 H, H-2"), 4.50 (t, J = 7.8 Hz, 1 H, H-1), 3.00 (m, 1 H, H-3), 2.60 (m, 1 H, H<sub>a</sub>-2), 2.40 (m, 1 H, H<sub>b</sub>-2), 1.38 (d, J = 7.0 Hz, 3 H, Me); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75.5 MHz)  $\delta$  136.45, 136.39, 136.37 (s each, C-7a',7a",7a"), 126.94 (s, C-3a"3a"), 126.64 (s, C-3a'), 122.29 (s, C-3'), 121.64, 121.62, 121.58 (d each, C-6',6"6"), 121.48 (d, C-2",2"), 120.22 (d, C-2'), 120.01 (s, C-3"), 119.96 (s, C-3"), 119.66 (d, C-4'), 119.57 (d, C-4", 4"), 118.87 (d, C-5"5"), 118.79 (d, C-5'), 111.13 (d, C-7"), 111.10 (d, C-7", 111.06 (d, C-7'), 43.59 (t, C-2), 31.78 (d, C-1), 28.80 (d, C-3), 21.81 (q, C-4); EIMS (70 eV) *m/z* (%) 403 (40), 286 (10), 258 (25), 245 (100), 217 (10), 145 (40), 117 (15); (-)-ESIMS *m/z* 402.8 [M-H<sup>+</sup>]; (+)-ESIMS *m/z* 426.7 [M+Na<sup>+</sup>]; HREIMS *m/z* 403.2030 (calcd. 403.20484 for C<sub>28</sub>H<sub>25</sub>N<sub>3</sub>).

**Biological Tests**. All tests were performed as described previously<sup>4</sup> using the agar diffusion method with bacteria (*Escherichia coli*, *Staphylococcus aureus*, *Bacillus subtilis* and *Streptomyces viridochromogenes*) and fungi (*Candida albicans*, *Mucor miehei*) as test organisms.

Acknowledgment. We would like to thank Mrs. F. Lissy for technical assistance, Dr. H. Frauendorf and Mr. R. Machinek for the spectral measurements. This work was financed by a grant from the Volkswagenstiftung within the Lower Saxony Cooperative Research Project of Marine Biotechnology.

#### **References and Notes**

- Art. No. XXIII on Marine Bacteria. For Art. XXII, see: Maskey, R. P.; Li, F.; Qin, S.; Fiebig, H. H.; Laatsch, H.: Chandrananimycins A ~ C: Production of Novel Anti-cancer Antibiotics from a Marine *Actinomadura* sp. Isolate M048 by Variation of Medium Composition and Growth conditions. *J. Antibiot.*, accepted 2003.
- (2) Bergman, J.; Tilstam, U. Tetrahedron, 1985, 41, 2883–2884.
- (3) Laatsch, H. AntiBase 2000, A Natural Products Database for Rapid Structure Determination. Chemical Concepts, Weinheim 2000 and annual updates; see Internet http://www.gwdg.de/~ucoc/Laatsch/

- (4) Biabani, M. A. F.; Baake, M.; Lovisetto, B.; Laatsch, H.; Helmke, E.; Weyland, H. J. *Antibiot.* **1998**, *51*, 333–340.
- (5) Maskey, R. P.; Asolkar, R. N.; Kapaum, E.; Wagner-Doubler. I.; Laatsch, H. J. Antibiot.
  2002, 55, 643–649.
- (6) Khuzhaev, V. U.; Aripova, S. F. Chem. Nat. Comp., 1998, 34, 108-109.
- (7) Pindur, U.; Müller, J. Chem. Ztg. 1986, 110, 333–335. Korolev, A.M.; Yudina, L. N.;
   Lazhko, L.I.; Reznikova, M.I.; Preobrazhenskaya, M. N. Chem. Heterocycl. Comp. 1999, 35, 561–569.
- (8) Pouchert, C. J; Campbell, J. R. *The Aldrich Library of 13C and 1H FT NMR Spectra*;
  Aldrich Chemical Co.: Milwaukee, **1992**; Vol. 3, 138A, 158B. Turner, W. B.; Aldridge,
  D. C.; Galt, S.; Giles D. *J. Chem. Soc.* (C) **1971**, 1623–1627.
- (9) Bell, R.; Carmeli, S.; Sar, N. J. Nat. Prod. 1994, 57, 1587-1590.
- (10) Kobayashi, M.; Aoki, S.; Gato, K.; Matsunami, K.; Kurosu, M.; Kitagawa, I. Chem.
   Pharm. Bull. 1994, 42, 2449–2451.
- (11) Shaaban, M.; Maskey, R. P.; Wagner-Dobler, I.; Laatsch, H. J. Nat. Prod. 2002, 65, 1660–1663.
- (12) Huang, Q.; Tezuka, Y.; Kikuchi, T.; Nishi, A.; Tubaki, K. Chem. Pharm. Bull. 1995, 43, 1035–1038.
- (13) Dembitsky, V. M.; Shkrob, I.; Lev, O. J. Chem. Ecol. 2000, 26, 1359–1366.
- (14) Zhungietu, G. I.; Chukhrii, F. N. Zh. Vses. Khim. Obshchest. 1970, 15, 228.
- (15) Gillespie, D. E.; Brady, S. F.; Bettermann, A. D.; Cianciotto, N. P.; Liles, M. R.; Rondon, M.R.; Clardy, J.; Goodman, R. M.; Handelsman, J. *Appl. Environ. Microb.* 2002, 68, 4301–4306.
- (16) Budzikiewicz, H.; Eckau, H.; Ehrenberg, M. Tetrahedron Lett. 1972, 36, 3807–3810.
- (17) Gao, X.; Petroff, B. K.; Oluola, O.; Georg, G.; Terranova, P. F.; Rozman, K. K. *Toxicol. Appl. Pharmacol.* **2002**, *183*, 179–188.
- (18) Hong, C.; Kim, H.-A.; Firestone, G. L.; Bjeldanes, L. F. *Carcinogenesis* 2002, 23, 1297–1305.

- (19) Bell, M.; Schmidt-Grimminger, D.-C. U.S. Patent **2002**, US 6399645; Chem. Abstr. 136:395952.
- (20) Ellenberger, S. R.; Ellenberger, W. P.; Andrus, G. M. 2001, WO Patent 0108637; Chem. Abstr. 134:141751.
- (21) Broadbent, T. A.; Broadbent, H. S. Curr. Med. Chem. 1998, 5, 337-352.
- (22) Jackson, A. H.; Prasitpan, N.; Shannon, P. V. R.; Tinker, A. C. J. Chem. Soc. Perkin Trans I 1987, 2543–2551.
- (23) Gil-Turnes, M. S.; Hay, M. E.; Fenical, W. Science 1989, 246, 116-118.
- (24) Kawasaki, W.; Matsui, K.; Akakabe, Y.; Itai, N.; Kajiwara, T. *Phytochemistry* **1998**, *47*, 27–29.
- (25) Maskey, R. P.; Asolkar, R. N.; Helmke, E.; Laatsch, H. J. Antibiot. 2002, 55, 893-898.





R

, R'







**4a**: R = H **4b**: R = Me



