

Elevated atmospheric CO₂ increases microbial growth rates in soil: results of three CO₂ enrichment experiments

EVGENIA BLAGODATSKAYA*†, SERGEY BLAGODATSKY†, MAXIM DORODNIKOV* and YAKOV KUZUYAKOV*

*Department of Agroecosystem Research, BayCEER, University of Bayreuth, 95440 Bayreuth, Germany, †Institute of Physicochemical and Biological Problems in Soil Science, Russian Academy of Sciences, 142290 Pushchino, Russia

Abstract

Increasing the belowground translocation of assimilated carbon by plants grown under elevated CO₂ can cause a shift in the structure and activity of the microbial community responsible for the turnover of organic matter in soil. We investigated the long-term effect of elevated CO₂ in the atmosphere on microbial biomass and specific growth rates in root-free and rhizosphere soil. The experiments were conducted under two free air carbon dioxide enrichment (FACE) systems: in Hohenheim and Braunschweig, as well as in the intensively managed forest mesocosm of the Biosphere 2 Laboratory (B2L) in Oracle, AZ. Specific microbial growth rates (μ) were determined using the substrate-induced respiration response after glucose and/or yeast extract addition to the soil. For B2L and both FACE systems, up to 58% higher μ were observed under elevated vs. ambient CO₂, depending on site, plant species and N fertilization. The μ -values increased linearly with atmospheric CO₂ concentration at all three sites. The effect of elevated CO₂ on rhizosphere microorganisms was plant dependent and increased for: *Brassica napus* = *Triticum aestivum* < *Beta vulgaris* < *Populus deltoides*. N deficiency affected microbial growth rates directly (N limitation) and indirectly (changing the quantity of fine roots). So, 50% decrease in N fertilization caused the overall increase or decrease of microbial growth rates depending on plant species. The μ -value increase was lower for microorganisms growing on yeast extract than for those growing on glucose, i.e. the effect of elevated CO₂ was smoothed on rich vs. simple substrate. So, the r/K strategies ratio can be better revealed by studying growth on simple (glucose) than on rich substrate mixtures (yeast extract). Our results clearly showed that the functional characteristics of the soil microbial community (i.e. specific growth rates) rather than total microbial biomass amount are sensitive to increased atmospheric CO₂. We conclude that the more abundant available organics released by roots at elevated CO₂ altered the ecological strategy of the soil microbial community specifically a shift to a higher contribution of fast-growing r-selected species was observed. These changes in functional structure of the soil microbial community may counterbalance higher C input into the soil under elevated atmospheric CO₂ concentration.

Keywords: *Beta vulgaris*, Biosphere 2, *Brassica napus*, elevated CO₂, FACE, glucose, microbial biomass, microbial growth rates, *Populus deltoides*, r and K strategy, rhizodeposition, rhizosphere microorganisms, substrate-induced respiration, *Triticum aestivum*

Received 6 March 2009; revised version received 1 June 2009 and accepted 4 June 2009

Introduction

The mineralization activity of soil microorganisms is crucial in regulating CO₂ release into the atmosphere

from soils, where about 60% of the carbon (C) stock of terrestrial ecosystems is sequestered. An increased CO₂ concentration in the atmosphere of up to 450–600 ppm, as predicted for 2050 (Houghton *et al.*, 2001), cannot directly affect soil mineralization processes because the CO₂ concentration in soil is several orders of magnitude higher. Nonetheless, indirect plant-mediated effects of

Correspondence: Sergey Blagodatsky, tel. +7 4967 736 367, fax +7 4967 330 595, e-mail: sblag@mail.ru

elevated CO₂ concentration – including more rhizodeposits and root exudates and changes in their composition (van Veen *et al.*, 1991; Diaz *et al.*, 1993; Hungate *et al.*, 1997; Cheng, 1999) – can affect the amount, composition and functions of soil microorganisms (Montealegre *et al.*, 2002; Carney *et al.*, 2007; Kandeler *et al.*, 2008). Surprisingly, recent studies have revealed only minor changes in soil microbial communities under elevated CO₂, especially in total prokaryotic and eukaryotic abundance (Lipson *et al.*, 2006; Haase *et al.*, 2008; Kanerva *et al.*, 2008; Lesaulnier *et al.*, 2008). Microbial biomass increased slightly (Ross *et al.*, 1995; Insam *et al.*, 1999, reviewed by De Graaff *et al.*, 2006) or even decreased (Hungate *et al.*, 1996) under elevated vs. ambient CO₂, and microbial respiration showed a more consistent increase – ca. 17% according to De Graaff *et al.* (2006). It remains unclear, however, which factors were responsible for the absence of strong effects of elevated CO₂ on microorganisms despite the greater quantity and availability of organic substrates released by roots (Zak *et al.*, 2000b; Freeman *et al.*, 2004). Increased input of available C by rhizodeposits does not activate SOM-mineralizing microorganisms (usually K-strategists): numerous long-term studies found no or a minor effect of elevated CO₂ on SOM content (Schlesinger & Lichter, 2001; Van Groenigen *et al.*, 2003; Lichter *et al.*, 2005; De Graaff *et al.*, 2006 and references therein). Rapid decomposition of increased C inputs by more active microbial biomass under elevated CO₂ can potentially explain the unaffected content of recalcitrant soil organic carbon under such CO₂ conditions (Lichter *et al.*, 2005; Taneva & Gonzalez-Meler, 2008). We therefore hypothesized that the increase of available C in the rhizosphere at elevated CO₂ will alter the structure and activity of the soil microbial community. The result is an increase of fast-growing r-strategists that quickly metabolize easily available substrates. In such a case, a greater input of available substrates by root exudates can be counterbalanced by faster microbial turnover, leading to higher rates of substrate mineralization.

Stimulation of fast-growing microorganisms by elevated CO₂ can be evaluated by examining the growth rates of the whole microbial community. Thus, *in situ* investigations of microbial growth rates are necessary to support the hypothesized increase of fast-growing microorganisms in the rhizosphere of different plant species under elevated CO₂. The lack of experimental studies relating specific microbial growth rates (μ) in soil with increasing atmospheric CO₂ is due to several difficulties: (1) direct measurements in pure cultures do not adequately reflect real rates of processes and microbial interactions in soil; and (2) until now there are no *direct* methods for satisfactorily estimating microbial growth *in situ*. An *indirect* approach suitable for estimat-

ing μ is based on the kinetics of substrate-induced respiration (KSIR) (Blagodatsky *et al.*, 2000; Stenström *et al.*, 2001). This approach characterizes the growth rates of whole microbial community according to microbial growth model proposed by Panikov (1995). The model reflects the transition of soil microorganisms from a 'sustaining' (Smith *et al.*, 1986) to an active state, considering both the lag-phase and phase of exponential growth after substrate addition. Beyond estimating μ , the KSIR approach also reveals the sustaining and growing fractions of microbial biomass (Panikov & Sizova, 1996; Blagodatsky *et al.*, 2000). If the available substrates in the rhizosphere increase under elevated CO₂, we expected more pronounced changes of microbial growth rates in the rhizosphere than in root-free soil.

Glucose is commonly used to induce the respiratory activity of soil heterotrophic microorganisms (Anderson & Domsch, 1978). Note that soil contains not only oligotrophic species capable of efficient growth on glucose, but also auxotrophic microorganisms, those that cannot grow in the absence of vitamins, amino acids, and growth factors (Alexander, 1991; Yadav, 2007). Accordingly, we applied – along with glucose and mineral salts – mixtures containing yeast extract in order to stimulate a broader spectrum of soil microorganisms with diverse nutrient requirements. Most of the C entering the soil as litter or rhizodeposits under elevated CO₂ consists of energy-rich, easily available carbon sources such as low molecular sugars, and dicarboxylic acids (Jones *et al.*, 2004) of poor quality, e.g. with a wide C:N ratio caused by lower N availability (Grayston *et al.*, 1998; Lesaulnier *et al.*, 2008). We therefore hypothesized lower nutrients requirements of microbial communities under elevated CO₂.

Soil N availability controls the increase of above- and belowground plant biomass against the background of elevated CO₂. Depending on the intensity of N limitation under such conditions the root growth may decrease (Pregitzer *et al.*, 2000) or, on the contrary, the root mass (especially that of fine roots) may increase (Hyvonen *et al.*, 2007). Thus, the mineralization activity of microorganisms under elevated CO₂ is dependent on both interactions with the plant community and with N availability (Freeman *et al.*, 2004). This causes the discrepant results and difficulties in investigating the effect of elevated CO₂ on microbial activity and mineralization processes in soil. Accordingly, the hypothesized shift in the soil microbial community from K- to r-strategists can be also affected by N availability.

We investigated the long-term effect of elevated CO₂ on microbial biomass and specific growth rates of microorganisms of root-free and rhizosphere soil in field experiments under three different CO₂ enrichment

systems at spatially remote locations. The following questions were investigated in detail: how does the effect of elevated CO₂ on microbial growth rates in soil depend on (1) the CO₂ concentration, (2) the plant species, and (3) N fertilization. In order to stimulate a broader spectrum of soil microorganisms, we investigated the shift in metabolic functioning of the soil microbial community using three organic-mineral mixtures of easily available substrates.

Materials and methods

Site descriptions and soils

Free air carbon dioxide enrichment (FACE) at Stuttgart-Hohenheim, Germany. The FACE facility in Hohenheim (FACE-H) is located south of Stuttgart, Baden-Württemberg, Germany (48°43'N, 9°13'E 100 m a.s.l.). The soil is a Gleyic Cambisol (WRB, 1998) without CaCO₃. The soil properties were identical under ambient and elevated CO₂ treatments (Table 1).

The CO₂ enrichment experiment, starting in 2002, included plots with elevated atmospheric CO₂ (540 ppm, with enclosures), ambient plots (380 ppm, with enclosures), and control plots (ambient CO₂ level, no enclosures) (Erbs & Fangmeier, 2006). Each treatment was replicated five times. Spring wheat (*T. aestivum* L. cv Triso) was annually planted on the plots from 2002 to 2006. In 2007, oilseed rape (*Brassica napus* L.) was grown on the plots for the first time. Beginning in 2003, inorganic NPK fertilizers were annually applied in equal amounts of 140 kg N ha⁻¹, 60 kg K ha⁻¹, 30 kg P ha⁻¹ to each plot under ambient and elevated CO₂ treatments. No organic fertilizers were applied. Soil was sampled from the top 10 cm of each ambient and elevated-CO₂ plot using soil corers (inner diameters: 5 cm) in September 2007, 3 weeks after the rape harvest. Soil samples were taken from locations that adhered to the plant roots and stored at 7 °C for 1 week before analyses.

FACE at the Institute of Agroecology, FAL, Braunschweig, Germany. The second FACE facility (FACE-B) used in the study was located at the experimental farm of the Federal Agricultural Research Centre (FAL) in Braunschweig, Lower Saxonia, Germany (52°18'N, 10°26'E, 79 m a.s.l.). Investigations were carried out on plots under ambient (350 ppm) and elevated (550 ppm) atmospheric CO₂ concentrations, values which had been applied since May 2000 (Weigel *et al.*, 2006). Two plant species, sugar beet (*Beta vulgaris* L.) and winter wheat (*T. aestivum* L.), grown in crop rotation were investigated. The soil type was loamy-sand Cambisol derived from loess (Table 1). N supply was restricted to

Table 1 Overview of the three CO₂ enrichment experiments and the relevant soil properties

Site location, acronym	Plant species	Atmospheric CO ₂ (ppm)		Soil type	C (%)	N (%)	Sand (%)	Silt (%)	Clay (%)	pH _{KCl}	Reference
		Ambient	Elevated								
Braunschweig FACE-B	<i>Beta vulgaris</i> , <i>Triticum aestivum</i>	350	550	Cambisol, loamy sand	1.1	0.09	42	52.6	5.4	6.8	Weigel & Dämmgen (2000)
Hohenheim FACE-H	<i>Brassica napus</i>	380	540	Gleyic Cambisol	1.59	0.17	9	69	22	6.8	Erbs & Fangmeier (2006)
Oracle, AZ, B2L	<i>Populus deltoides</i>	400	800 or 1200	Silty loam	2	0.19	28	55	17	7.2	Torbert & Johnson (2001)

FACE, free air carbon dioxide enrichment; B2L, Biosphere 2 Laboratory.

50% of adequate N in half of each of the FACE-B plots, and the soils were annually amended with mineral fertilizers at two different rates: 126 and 63 kg N ha⁻¹ yr⁻¹. Soils were sampled after 3 and 4 years from the beginning of the fumigation with elevated CO₂.

Composite soil samples were taken in three replicates from the layer 0 to 10 cm during harvesting time (sugar beet) or 1 week after cropping the winter wheat. Rhizosphere soil was taken from locations that adhered to the plant roots, whereas root-free soil was taken from the space between rows of crops. Soil was stored field-fresh in aerated polyethylene bags for a maximum 8 weeks at 4°C. Samples taken from this experiment allow estimating the rhizosphere and N amendment effects along with the effect of elevated CO₂ on the SIR kinetics.

Intensively managed forest mesocosm, 'Biosphere 2,' in Oracle, AZ, USA. The Biosphere 2 Laboratory (B2L) is 1.27 ha virtually airtight environmental research facility located in Oracle, AZ (32°34'N, 110°51'W; 1200 m a.s.l.). Soil samples were taken from experimental plots belonging to the intensively managed forest mesocosm (Griffin *et al.*, 2002) with eastern cottonwood (*Populus deltoides* Bartr. ex Marsh.) grown under ambient (400 ppm) and under two levels of elevated (800 and 1200 ppm) atmospheric CO₂ concentrations. The design and operation of B2L are described in detail elsewhere (Zabel *et al.*, 1999). The soil is a silt loam (Torbert & Johnson, 2001), not fertilized, with properties presented in Table 1. Rhizosphere soil adhering to the plant roots was taken in three replicates (i.e. composite samples from three different trees) during root sampling, whereas root-free soil was taken in three replicates from the space under the trees, but >3–5 cm away from the root surface.

Not later than 1 day after sampling, the soils were sieved (<2 mm), fine roots and other plant debris were carefully removed, and the soils were stored at 4°C during 1–3 months. Before the analyses, soil moistures were adjusted to 60% of water holding capacity (WHC) and samples were preincubated for 24 h at 22°C.

Microbial biomass and the kinetics of substrate-induced respiration

Parameters of microbial growth were determined according to the dynamics of the CO₂ emission rate from soil amended with glucose and/or yeast extract. Mixtures containing yeast extract were applied to soil in order to stimulate a broader spectrum of soil microorganisms and to estimate the metabolic activity of microbial guilds other than glucose-responding heterotrophs.

Samples of 10 g (dry weight) soil were amended with a powder-mixture (I) containing 10 mg g⁻¹ glucose, 20 mg g⁻¹ talcum, and mineral salts: (NH₄)₂SO₄ – 1.9 mg g⁻¹, K₂HPO₄ – 2.25 mg g⁻¹, and MgSO₄·7H₂O – 3.8 mg g⁻¹. Mixtures II and III were prepared as in mixture I with the following modifications: 0.3 mg g⁻¹ of yeast extract LD (Difco Manual, 1998) was added to glucose (10 mg g⁻¹) in mixture II as a combined C source; 4 mg g⁻¹ of yeast extract was used in mixture III instead of glucose. Optimal concentrations of substrates were found in preliminary experiments, so that the soil amendments were sufficient to enable unlimited exponential growth of soil microorganisms during the initial period. Mineral salts included in mixtures were selected so that the substrate changed the pH of soil for <0.1.

The CO₂ production rate was measured hourly at 22°C using an automated infrared-gas analyzer system (Heinemeyer *et al.*, 1989).

Soil microbial biomass C (C_{mic}) was determined using the initial rate of substrate-induced respiration when soil was amended with mixture I according to the equation of Anderson & Domsch (1978):

$$C_{mic}(\mu\text{g g}^{-1} \text{ soil}) = (\mu\text{L CO}_2 \text{ g}^{-1} \text{ soil h}^{-1}) \times 40.04. \quad (1)$$

Specific growth rate (μ) was determined by fitting of equation parameters to the experimental data on CO₂ evolution rate (v) according to:

$$v(t) = A + B \times \exp(\mu \times t), \quad (2)$$

where A is the initial rate of uncouple (nongrowth) respiration, B is the initial rate of couple (growth) respiration, t is the time (Blagodatsky *et al.*, 2000; Panikov & Sizova, 1996). The parameters of Eqn (2) were determined by fitting MODELMAKER © Version 3.0.3 software (Modelmaker, 1997). Fitting was restricted to the part of the curve that corresponded better to applied model Eqn (2), as indicated by maximum values of statistic criteria: r^2 , the fraction of total variation explained by the model defined as the ratio of the model weighted sum of squares to the total weighted sum of squares.

Statistical analysis

All data presented are mean values from three replications. Nonparametric tests (Mann–Whitney and Wilcoxon) were used for estimating the significant effects ($P < 0.05$) of the following factors: ambient vs. elevated CO₂ (CO₂); root adhered vs. root-free soil (Rhi); and different N amendment (N). Three-way analysis of variance (ANOVA) was applied to split the effects of the three mentioned factors and their interactions for two blocks of data: soil under sugar beet or under winter

wheat (FACE-B). Such analysis was appropriate only for FACE-B experiment where effects of three factors could be measured without interference of time and plant species. Two-way completely randomized ANOVA was applied for FACE-B experiment in order to characterize effects of elevated CO₂ and grown crop. When significant effects were found a multiple comparison using Duncan's multiple range test (significance level 5%) were performed. Regression analysis was used to check the effect of CO₂ concentration on microbial specific growth rates. All variables treated passed normality and equal variance tests.

Results

Microbial biomass and KSIR under elevated CO₂

Microbial biomass ranged from 204 to 429 µg C g soil⁻¹ under ambient CO₂ concentrations (Table 2). Significantly larger microbial biomass C was measured in FACE-H soil compared with that in B2L and FACE-B soils. No significant effect of elevated CO₂ on total microbial biomass C was observed in any soil if direct comparison of means or two-way ANOVA was applied, although in all cases the microbial biomass at elevated CO₂ tended to increase (Table 2). If influence of all three studied factors (atmospheric CO₂, plant roots, N application rate) is considered for FACE-B experiment, then elevated CO₂ effect on total microbial biomass under sugar beet is significant at $P < 0.05$ level according to three-way ANOVA (Table 3).

For all soils the increase in the respiratory response after glucose addition was steeper under elevated vs. ambient CO₂ (Fig. 1). The steepness of the respira-

tion curves increased considerably with increasing CO₂ concentration (Fig. 1, bottom right). For each soil type, specific microbial growth rates (μ) estimated by Eqn (2) were significantly higher under elevated CO₂ (Table 2). Regression analysis clearly revealed significant positive relationships between increasing atmospheric CO₂ concentration and μ -values for all soils and treatments (Fig. 2).

Plant species and root vicinity effect on microbial growth rates

Significant differences ($P < 0.05$) in specific microbial growth rates were observed between the three experimental sites for soils under ambient CO₂. Within the same FACE system and the same soil type (FACE-B), the differences between μ -values under different plant species *B. vulgaris* (root crop) and *T. aestivum* (cereal) were insignificant at ambient CO₂ concentration (Table 2). Thus, under ambient CO₂ the effect of soil type on μ -values was higher than the effect of plant species. The same tendency was also observed under elevated CO₂ (Table 2).

In root-free soil as well as in the rhizosphere of all plant species, the μ -values were always higher under elevated than under ambient CO₂ (Fig. 3), i.e. all ratios presented in Fig. 3 are higher than 1.0. The effect of elevated CO₂ on μ in the rhizosphere of different plants increased in the following order: *B. napus* (540 ppm) = *T. aestivum* (550 ppm) < *B. vulgaris* (550 ppm) < *P. deltooides* (800 ppm) (Fig. 3). In the last case, both the plant type and higher CO₂ concentration likely affected the microbial response. The effect of elevated CO₂ on μ tended to be more pronounced in the rhizosphere than

Table 2 Soil microbial biomass, and specific growth rates (μ) of rhizosphere microorganisms in the three CO₂ enrichment experiments

Site location	Plant species	Atmospheric CO ₂ (ppm)	Microbial biomass, (µg C × g ⁻¹)	μ (h ⁻¹)
Braunschweig, FACE-B*	<i>Beta vulgaris</i>	350	303 Ab	0.246 Aa
		550	337 Ab	0.284 Ba
	<i>Triticum aestivum</i>	350	204 Aa	0.255 Aa
		550	206 Aa	0.297 Ba
Hohenheim, FACE-H*	<i>Brassica napus</i>	380	429 A	0.194 A
		540	464 A	0.207 B
Oracle, AZ, B2L†	<i>Populus deltooides</i>	400	296 A	0.301 A
		800	385 BC	0.393 B
		1200	322 AB	0.475 C

Treatments followed by the same letters are not significantly different between elevated and ambient CO₂ (uppercase letters) and various plants (FACE-B, lowercase letters) at $P \leq 0.05$.

*Data for rhizosphere soil with full rate of N application.

†Data for rhizosphere soil.

FACE, free air carbon dioxide enrichment; B2L, Biosphere 2 Laboratory.

Table 3 Input (%) of atmospheric CO₂ level (CO₂), N application rate (N), and plant roots (Rhi) into variation of microbial specific growth rates (μ) and total biomass estimated by three-way ANOVA for FACE-B experiment

Crop	Source of variation	Specific growth rate (μ)	Total microbial biomass
<i>Beta vulgaris</i>	CO ₂	58*	28*
	Rhi	12*	48*
	N	16*	10*
	CO ₂ × Rhi	6*	1
	CO ₂ × N	0	1
	Rhi × N	1	0
	CO ₂ × Rhi × N	4*	3
	Unexplained	4	11
<i>Triticum aestivum</i>	CO ₂	35*	4
	Rhi	0	4
	N	25*	2
	CO ₂ × Rhi	0	0
	CO ₂ × N	0	0
	Rhi × N	3	28*
	CO ₂ × Rhi × N	18*	5
	Unexplained	17	56

*Input significant at $P \leq 0.05$.

FACE, free air carbon dioxide enrichment; ANOVA, analysis of variance.




Fig. 1 Kinetics of substrate-induced respiration for soil under three plant species grown at different levels of atmospheric CO₂ in three different CO₂ enrichment experiments. Measured data are presented as symbols, simulation results as lines [Eqn (2)].

in the soil sampled between the rows. This tendency was significant only for soils under *B. vulgaris* (FACE-B) and under *P. deltoides* (B2L) at highest CO₂ concentration (Fig. 3).

Combined effect of elevated CO₂ and nitrogen fertilization on microbial growth rates

Both at ambient and at elevated CO₂ concentrations, the μ -values at plots with *B. vulgaris* tended to be higher at

half vs. the full rate of N fertilization (Fig. 4, top, FACE-B). The opposite effect was observed in soils under *T. aestivum* (FACE-B, data not shown), but in general, N fertilization significantly affects specific growth rates under both cultures as is confirmed by three-way ANOVA results for FACE-B experiment (Table 3). Under *B. vulgaris* at half N rate, the effect of elevated CO₂ on μ -values (presented as ratio of μ_{elevated} to μ_{ambient}) was significantly higher in rhizosphere than in root-free soil (Fig. 4, bottom, FACE-B). However, there were no

differences in the elevated CO₂ effect between rhizosphere and root-free soil at the full rate of N fertilizers (Fig. 4, bottom). The 2 × 2 × 2 ANOVA (FACE-B data) showed no significant ($P < 0.05$) interactions between paired factors affecting the μ and total microbial biomass, i.e. CO₂ × N and Rhi × N for both crops. The only exception was the μ increase for CO₂ × Rhi interaction under sugar beet. In latter case significant rhizosphere effect in respect to μ was enhanced by elevated CO₂. Three factors interaction CO₂ × Rhi × N influences significantly specific growth rates under both plant species.

Effect of elevated CO₂ on microorganisms with different nutrient requirements

The comparison of microbial growth on glucose and on yeast extract was used to estimate the response to elevated




Fig. 2 Specific growth rate of microorganisms on glucose at different levels of atmospheric CO₂. Data for three experimental sites are shown.




Fig. 3 Relative increase of microbial specific growth rate (μ) caused by CO₂ enrichment in root-free and rhizosphere soil of three CO₂ enrichment experiments. Bars show standard errors, $n = 3$.

CO₂ of soil microorganisms with different nutrient requirements (FACE-B experiment). The lag-period before the start of exponential growth was 4–9 h shorter (Table 4) and the increase in respiration rate was steeper (Fig. 5, top) for microorganisms grown on yeast extract vs. glucose both under ambient and elevated CO₂. The maximal specific-growth rate was 2.1–2.7-fold higher on yeast extract (Table 4). Thus, microorganisms utilizing yeast extract differ (at least partly) from those metabolizing glucose in their metabolic abilities. Applying yeast extract along with glucose helps to estimate the effects of elevated CO₂ on a more diverse metabolic spectrum of soil microorganisms.

The differences in steepness of microbial respiration curves at ambient and elevated CO₂ (Fig. 5, top) were maximal at growth on mixture I (glucose with mineral salts) and were less pronounced by increasing the amount of added yeast (mixtures II and III). Similarly to glucose (Fig. 1, Table 2), the growth on yeast extract had significantly higher μ -values at elevated vs. ambient CO₂ (Table 4). However, the increase at elevated (vs. ambient) CO₂ was stronger for growth on glucose than on yeast extract (Fig. 5, bottom). This shows the lower effect of elevated CO₂ on auxotrophic microorganisms compared with microorganisms capable of growth on glucose with mineral nutrients.

Discussion

Effect of elevated CO₂ on microbial growth rates

The microbial growth rates were significantly higher under elevated vs. ambient CO₂ in the three




Fig. 4 Effect of nitrogen fertilization on microbial specific growth rate μ (top) and its relative increase (bottom) under sugar beet (*Beta vulgaris*) grown at ambient and elevated CO₂ in free air carbon dioxide enrichment-B experiment. Bars shows standard errors, $n = 3$.

Table 4 Specific growth rates (μ) of microbial community growing on yeast extract and index of auxotrophy for FACE experiment in Braunschweig

Crop	Atmospheric CO ₂ (ppm)	μ_{\max} (h ⁻¹)	Lag-time (h)	Index of auxotrophy (μ) glucose/ μ yeast extract
<i>Beta vulgaris</i>	350	0.608 A	6.1 A	0.405 A
	550	0.637 B	6.2 A	0.446 B
<i>Triticum aestivum</i>	350	0.588 A	6.4 B	0.434 A
	550	0.64 B	5.7 A	0.464 A

Treatments followed by the same letters are not significantly different between elevated and ambient CO₂ at $P \leq 0.05$. FACE, free air carbon dioxide enrichment.

investigated CO₂ enrichment systems, for every studied soil type, plant species and N fertilization rate. According to the ecological theory of life strategies, under nonlimiting conditions, microorganisms with r-strategy are characterized by higher μ -values and less efficient

substrate utilization than K-strategists (Andrews & Harris, 1986). Hence, plants growing in an elevated CO₂ atmosphere selectively increased the fraction of r-strategists in soil microbial communities. A strong increase in CO₂ concentration (from 400 to 1200 ppm in




Fig. 5 CO₂ evolution rate (top) and relative increase of microbial specific growth rate μ (bottom) for free air carbon dioxide enrichment-B soil under sugar beet (root-free, full N application rate) after different substrate addition: odd numbers (filled symbols) – plots under elevated atmospheric CO₂, even numbers (empty symbols) – plots under ambient CO₂. Substrates added: 1, 2 – glucose (mixture I); 3, 4 – glucose + yeast extract (mixture II); 5, 6 – yeast extract (mixture III). Treatments followed by the different letters are significantly different at $P < 0.05$.

B2L experiment) yielded a 58% increase in μ -values. However, at the smaller increase (from 350 to 550 ppm in both FACE experiments) the μ -values increased by 0.013–0.038 h⁻¹, i.e. 7–15% of the μ -values under ambient CO₂. Even such relatively slight increases in growth rates can have important consequences because exceedingly small differences in the specific growth rates (as small as 0.0055 h⁻¹) of two populations may be sufficient for the marginally faster growing organisms to be more competitive (Mason & Slater, 1979). Hodge *et al.* (1998) reported more cultivated bacteria (usually r-strategists) as along with more intensive rates of substrate utilization on BIOLOG-plates in soil with ryegrass grown under elevated vs. ambient CO₂ concentrations. These observations were supported by quantitative-PCR analyses that revealed more heterotrophic decomposers within a bacterial community at elevated CO₂ (Lesaulnier *et al.*, 2008). However, total bacterial and eukaryotic abundance as well as the

abundance of denitrifying bacteria were not affected by elevated CO₂ (Haase *et al.*, 2008; Lesaulnier *et al.*, 2008). We suggest that the functional characteristics of soil microbial community (microbial growth or substrate mineralization rates) rather than total microbial biomass are sensitive to increased atmospheric CO₂.

Most studies using relatively conservative indexes of soil microbial communities – DNA content, total PLFA and diversity estimated based on PLFA analysis – registered no or only a subtle effect of elevated CO₂ (Griffiths *et al.*, 1998; Zak *et al.*, 2000b; Ebersberger *et al.*, 2004; Lipson *et al.*, 2006; Denef *et al.*, 2007; Kanerva *et al.*, 2008; Paterson *et al.*, 2008). The studies using community level physiological profiling (CLPP) and similar approaches (Rillig *et al.*, 1997; Grayston *et al.*, 1998; Mayr *et al.*, 1999), showed, on the contrary, distinct response of microbial community to elevated CO₂. Direct comparison of total (DNA-based) and active (RNA-based) bacterial communities (Jossi *et al.*, 2006) showed that p CO₂ increase mainly influenced active and root-associated bacterial components. After adding glucose, potential enzyme activities under elevated CO₂ were up to twofold higher than under ambient CO₂ (Dorodnikov *et al.*, 2009b). Supporting our observations, this indicates the increased activity of fast-growing microorganisms, which leads to accelerated turnover of available carbon in soil under elevated CO₂. The measurement of specific growth rates of microorganisms responding to substrate additions can therefore serve as robust and simple method to quantitatively estimate changes in whole microbial community caused by environmental changes (Monson *et al.*, 2006; Lipson *et al.*, 2009) including p CO₂ increase (Dorodnikov *et al.*, 2009a).

The inconsistency of effects of elevated CO₂ on microbial biomass and activity observed in different studies (Zak *et al.*, 2000b) can be related to interactions with other factors such as nutrient limitation and the effect of plant species.

Rhizosphere and plant species effects

Elevated CO₂ increases microbial specific growth rates both in rhizosphere and in bulk soil (Fig. 3). Though the rhizosphere effect was observed for the μ response to elevated CO₂ in several cases, the altered functional properties of microbial communities included the whole soil volume and were not merely transient or linked with the current input of root exudates.

We find no direct effect of plant species (*B. vulgaris* and *Triticum aestivum*) on specific microbial growth rates within the same FACE system (FAL). This agreed with Deiglmayr *et al.* (2004), who found no differences between *Lolium perenne* L. and *Trifolium repens* L. effects on the nitrate-reducing community in the rhizosphere

under long-term elevated CO₂. Similarly to our observations, inconsistent effects of different plant species on soil microbial populations were found in the rhizosphere of *L. perenne* and of *T. repens* (Schortemeyer *et al.*, 1996; Montealegre *et al.*, 2002). Such inconsistency can be due to nonuniform temporal and spatial responses of root systems. According to Haase *et al.* (2007), the C flux from plant roots of *Phaseolus vulgaris* L. was significantly increased only in apical root zones. Moreover, even in those zones, the elevated CO₂ effect was inconsistent and resulted in an increase or decrease in rhizodeposits within the plant development stages (Haase *et al.*, 2008). Study of plant and soil lipid modification under elevated CO₂ argued against plant/soil turnover determinations for individual compounds (Wiesenberg *et al.*, 2008). Thus, studies focused on the effects of root exudations on microbial growth rates are necessary to reveal the effect of higher CO₂ levels under different plant species during the vegetation season.

Combined effect of nitrogen and CO₂ enrichment

Addition of N stimulates plant and microbial growth, and these effects may be modified by elevated CO₂ (Hungate *et al.*, 1996). Strongly increased plant growth by N fertilization may also increase the total amount of roots and rhizodeposits. However, this effect is frequently counterbalanced by a decreased percentage of C allocated belowground at higher N availability (Kuz'yakov & Domanski, 2000; Kuz'yakov *et al.*, 2001; Henry *et al.*, 2005).

Higher microbial specific growth rates under plots with 50% N supply (*B. vulgaris*, FACE-B, Blagodatsky *et al.*, 2006) coincide with larger fine root production in these conditions (Weigel *et al.*, 2006). Presumably, a trade-off in N uptake between plants and microbes took place in the rhizosphere of *B. vulgaris* (reduced N fertilization), leading to differences in specific growth rates of microbial communities under ambient and elevated CO₂. Thus, N enrichment smoothed the effect of elevated CO₂ on μ -values in the rhizosphere soil under *B. vulgaris*.

The significant effect of CO₂ × Rhi × N interactions along with the direct N effect on microbial specific growth rates (Table 3) supports the two-way influence hypotheses: N deficiency can affect microbial growth rates directly (by decreasing C use efficiency, Blagodatskiy *et al.*, 1993) and indirectly (through changing the quantity of fine roots). In the latter case plant species effect and root vicinity (rhizosphere effect) modify the effect of N fertilization. Thus, the effect of elevated CO₂ (higher μ -values) can be counterbalanced by N amendment. Similar to our observations, Chung *et al.* (2007) reported that the response of microbial communities to

elevated CO₂ was counterbalanced by N deposition in a study on the effect of plant species diversity on microbial biomass and fungal abundance in a BioCON-FACE experiment in east-central Minnesota. However, in another FACE experiment with *Populus*, N amendment stimulated C utilization rates estimated by the Microresp-CLPP method (Lagomarsino *et al.*, 2007). A 30% increase in soil heterotrophic respiration caused by elevated CO₂ in the same FACE experiment was not affected by N fertilization (Lagomarsino *et al.*, 2009), so the real availability of C and N in this case does not correspond to microbial growth rates obtained by the CLPP approach. Increased N availability reduced microbial respiration in a pine forest under elevated CO₂ at the Duke Free Air Carbon Enrichment site (Billings & Ziegler, 2008). These changes in total heterotrophic activity were apparently connected with the shift in microbial community structure estimated by $\delta^{13}\text{C}$ -PLFA analysis. Interestingly, Gram-positive actinomycetes, typical soil bacteria with features of K-strategists, i.e. lower maximal-specific growth rates, were abundant under elevated CO₂ and full N fertilization (Billings & Ziegler, 2008). Such findings agree well with our results showing lower μ under full N supply and elevated CO₂.

Thus, our study confirms a plant species effect on microbial response to N availability under elevated CO₂ (Hungate *et al.*, 1996). Contradictory N effects on microbial activity under different plant communities can be explained by the different N requirements of dominant microbial species in the rhizosphere of various plant species. Different effects of N fertilizers on root biomass as well as on the rhizodeposition of various plant species (Kuz'yakov *et al.*, 2001, 2002) are additional potential explanations for inconsistent interactive N × elevated CO₂ effects on microbial growth parameters.

Effect of elevated CO₂ on auxotrophic microorganisms

Applying three different organic-mineral mixtures to estimate microbial growth better mirrors metabolic diversity and the auxotrophic requirements of soil microorganisms than applying glucose only. The higher μ -values during growth on yeast extract vs. glucose indicated that yeast extract offers selective advantages to microorganisms that are able to grow rapidly on rich media but require growth factors. The greater the increase in μ -values on yeast extract vs. glucose, the greater the auxotrophic requirements of the microbial community. Thus, the ratio between maximal specific growth rates on glucose/yeast extract ($\mu_{\text{gl}}/\mu_{\text{ye}}$) may serve as an auxotrophy index of the soil microbial community (Blagodatskaya *et al.*, 2003). High auxotrophy indices correspond to communities with low auxotrophic requirements. For both crops studied in

the FACE-B experiment (*B. vulgaris* and *T. aestivum*), this index was higher under elevated vs. ambient CO₂ (Table 4). Thus, the portion of the microbial community lacking growth factors and vitamins seems to be lower in soils under elevated vs. ambient CO₂. Lower auxotrophic requirements at higher CO₂ concentrations can be due to an increased C input in the form of rhizodeposits with a higher C/N ratio (Cotrufo *et al.*, 1994, 2005). This results in the selection of microorganisms capable of rapid growth on energy-rich, simple substrates (Freeman *et al.*, 2004). This observation was indirectly confirmed by a strong decrease in the diversity of fungal species as well as of nitrate-reducing bacteria and archaea under elevated CO₂ (560 ppm), as estimated by ribosomal DNA sequences (Lesaulnier *et al.*, 2008).

The microbial respiration response induced by glucose was more sensitive to changes in atmospheric CO₂ concentration than the response induced by yeast extract (Fig. 5). This result agrees with Lipson *et al.* (2006), who also found glucose-induced SIR to be the best parameter reflecting the effect of elevated CO₂. We suppose that higher CO₂ levels stimulate rhizodeposition with a more uniform composition and high carbohydrate content; this benefits the development of r-strategists, whose respiration response is more distinct upon glucose amendment. Thus, the shift in specific growth rates reflecting the ratio between r- and K-strategists depends on the choice of test substrate. This r-K ratio can be better revealed by studying growth on simple than on rich substrate mixtures.

Conclusions

The significant increase of specific microbial growth rates was observed in three CO₂ enrichment experiments with different soils, plants, CO₂ levels, and N fertilization. This shift was probably mediated by increased rhizodeposition under elevated CO₂, resulting in a higher input of easily available substrates into the rhizosphere. N fertilization affects microbial growth rates both directly and through the modification of root biomass and thus, exudation amount. This bidirectional N influence may stimulate or counterbalance effect of elevated CO₂ depending on plant type, vegetation stage and general nutrient limitation. Thus, the functional characteristics of the soil microbial community (microbial growth and substrate mineralization rates) – rather than total microbial biomass amount – are sensitive to higher CO₂ concentrations in the atmosphere.

Elevated CO₂ decreased metabolic diversity and auxotrophy requirements of the soil microbial community. The higher (20 to 60%) specific growth rates reflect faster turnover of soil microorganisms under elevated CO₂. We therefore explain the absence of soil organic C and

microbial biomass accumulation in soil under elevated CO₂ (as might be expected because of higher C input into the rhizosphere) by faster C mineralization, confirmed by the faster turnover rates of soil microorganisms.

Acknowledgements

This research was supported by the European Commission (Marie Curie IIF grant for E. B., project MICROSOM) and by the Alexander von Humboldt Foundation (fellowship for S. B.). We are greatly thankful to the people, who made available soil samples from their CO₂ enrichment experiments: Prof. Dr A. Fangmeier (Hohenheim), Prof. Dr H.-J. Weigel and Dr T.-H. Anderson (Braunschweig), Prof. Karl Bil' and Prof. Barry Osmond (Biosphere 2 Laboratory).

References

- Alexander M (1991) Soil microbiology in the next 75 years: fixed, flexible, or mutable? *Soil Science*, **151**, 35–40.
- Anderson JPE, Domsch KH (1978) A physiological method for the quantitative measurement of microbial biomass in soils. *Soil Biology and Biochemistry*, **20**, 107–114.
- Andrews JH, Harris RF (1986) r and K-selection and microbial ecology. In: *Advances in Microbial Ecology*, **9** (ed. Marshall KC), pp. 99–144. Plenum Press, New York.
- Billings SA, Ziegler SE (2008) Altered patterns of soil carbon substrate usage and heterotrophic respiration in a pine forest with elevated CO₂ and N fertilization. *Global Change Biology*, **14**, 1025–1036.
- Blagodatskaya EV, Khokhlova OS, Anderson T-H, Blagodatskii SA (2003) Extractable Microbial DNA Pool and Microbial Activity in Paleosols of Southern Urals. *Microbiology*, **72**, 750–755.
- Blagodatskiy SA, Larionova AA, Yevdokimov IV (1993) Effect of mineral nitrogen on the respiration rate and growth efficiency of soil microorganisms. *Eurasian Soil Science*, **25**, 85–95.
- Blagodatsky SA, Blagodatskaya EV, Anderson TH, Weigel HJ (2006) Kinetics of the respiratory response of the soil and rhizosphere microbial communities in a field experiment with an elevated concentration of atmospheric CO₂. *Eurasian Soil Science*, **39**, 290–297.
- Blagodatsky SA, Heinemeyer O, Richter J (2000) Estimating the active and total soil microbial biomass by kinetic respiration analysis. *Biology and Fertility of Soils*, **32**, 73–81.
- Carney KM, Hungate BA, Drake BG, Megonigal JP (2007) Altered soil microbial community at elevated CO₂ leads to loss of soil carbon. *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 4990–4995.
- Cheng WX (1999) Rhizosphere feedbacks in elevated CO₂. *Tree Physiology*, **19**, 313–320.
- Chung H, Zak D, Reich P, Ellsworth D (2007) Plant species richness, elevated CO₂, and atmospheric nitrogen deposition alter soil microbial community composition and function. *Global Change Biology*, **13**, 980–989.
- Cotrufo MF, De Angelis P, Polle A (2005) Leaf litter production and decomposition in a poplar short-rotation coppice exposed to free air CO₂ enrichment (POPFACE). *Global Change Biology*, **11**, 971–982.

- Cotrufo MF, Ineson P, Rowland AP (1994) Decomposition of tree leaf litters grown under elevated CO₂: effect of litter quality. *Plant and Soil*, **163**, 121–130.
- De Graaff M-A, Van Groenigen K-J, Six J, Hungate B, Van Kessel C (2006) Interactions between plant growth and soil nutrient cycling under elevated CO₂: a meta-analysis. *Global Change Biology*, **12**, 2077–2091.
- Deiglmayr K, Philippot L, Hartwig UA, Kandeler E (2004) Structure and activity of the nitrate-reducing community in the rhizosphere of *Lolium perenne* and *Trifolium repens* under long-term elevated atmospheric pCO₂. *FEMS Microbiology Ecology*, **49**, 445–454.
- Denef K, Bubenheim H, Lenhart K, Vermeulen J, Van Cleemput O, Boeckx P, Müller C (2007) Community shifts and carbon translocation within metabolically-active rhizosphere microorganisms in grasslands under elevated CO₂. *Biogeosciences*, **4**, 769–779.
- Diaz S, Grime JP, Harris J, McPherson E (1993) Evidence of a feedback mechanism limiting plant response to elevated carbon dioxide. *Nature*, **364**, 616–617.
- Difco. (1998) Yeast extract. In: *Difco Manual*, 11th edn, pp. 572–574. Difco Laboratories, Sparks, MD, USA.
- Dorodnikov M, Blagodatskaya E, Blagodatsky S, Fangmeier A, Kuzyakov Y (2009a) Stimulation of r- vs. K-selected microorganisms by elevated atmospheric CO₂ depends on soil aggregate size. *FEMS Microbiology Ecology*, **69**, 43–52.
- Dorodnikov M, Blagodatskaya E, Blagodatsky S, Marhan S, Fangmeier A, Kuzyakov Y (2009b) Stimulation of microbial extracellular enzyme activities by elevated CO₂ depends on soil aggregate size. *Global Change Biology*, **15**, 1603–1614.
- Ebersberger D, Wermbter N, Niklaus PA, Kandeler E (2004) Effects of long term CO₂ enrichment on microbial community structure in calcareous grassland. *Plant and Soil*, **264**, 313–323.
- Erbs M, Fangmeier A (2006) Atmospheric carbon dioxide enrichment effects on ecosystems – experiments and real world. *Progress in Botany*, **67**, 441–459.
- Freeman C, Kim SY, Lee SH, Kang H (2004) Effects of elevated atmospheric CO₂ concentrations on soil microorganisms. *Journal of Microbiology*, **42**, 267–277.
- Grayston SJ, Campbell CD, Lutze JL, Gifford RM (1998) Impact of elevated CO₂ on the metabolic diversity of microbial communities in N-limited grass swards. *Plant and Soil*, **203**, 289–300.
- Griffin KL, Turnbull M, Murthy R *et al.* (2002) Leaf respiration is differentially affected by leaf vs. stand-level night-time warming. *Global Change Biology*, **8**, 479–485.
- Griffiths BS, Ritz K, Ebbelwhite N, Paterson E, Killham K (1998) Ryegrass rhizosphere microbial community structure under elevated carbon dioxide concentrations, with observations on wheat rhizosphere. *Soil Biology and Biochemistry*, **30**, 315–321.
- Haase S, Neumann G, Kania A, Kuzyakov Y, Römheld V, Kandeler E (2007) Elevation of atmospheric CO₂ and N-nutritional status modify nodulation, nodule carbon supply, and root exudation of *Phaseolus vulgaris* L. *Soil Biology and Biochemistry*, **39**, 2208–2221.
- Haase S, Philippot L, Neumann G, Marhan S, Kandeler E (2008) Local response of bacterial densities and enzyme activities to elevated atmospheric CO₂ and different N supply in the rhizosphere of *Phaseolus vulgaris* L. *Soil Biology and Biochemistry*, **40**, 1225–1234.
- Heinemeyer O, Insam H, Kaiser E-A, Walenzik G (1989) Soil microbial biomass and respiration measurements: an automated technique based on infra-red gas analysis. *Plant and Soil*, **116**, 191–195.
- Henry F, Nguyen C, Paterson E, Sim A, Robin C (2005) How does nitrogen availability alter rhizodeposition in *Lolium multiflorum* Lam. During vegetative growth? *Plant and Soil*, **269**, 181–191.
- Hodge A, Paterson E, Grayston SJ, Campbell CD, Ord BG, Killham K (1998) Characterisation and microbial utilisation of exudate material from the rhizosphere of *Lolium perenne* grown under CO₂ enrichment. *Soil Biology and Biochemistry*, **30**, 1033–1043.
- Houghton JT, Ding Y, Griggs DJ, Noguer M, Van Der Linden PJ, Xiaosu D (eds) (2001) *Climate Change 2001: The Scientific Basis Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change (IPCC)*. Cambridge University Press, Cambridge, UK, 944 pp.
- Hungate BA, Canadell J, Chapin FS III (1996) Plant species mediate change in soil microbial N in response to elevated CO₂. *Ecology*, **77**, 2505–2515.
- Hungate BA, Holland EA, Jackson RB, Chapin FS III, Mooney HA, Field CB (1997) The fate of carbon in grasslands under carbon dioxide enrichment. *Nature*, **388**, 576–579.
- Hyvonen R, Agren GI, Linder S *et al.* (2007) The likely impact of elevated [CO₂], nitrogen deposition, increased temperature and management on carbon sequestration in temperate and boreal forest ecosystems: a literature review. *New Phytologist*, **173**, 463–480.
- Insam H, Bååth E, Berreck M *et al.* (1999) Responses of the soil microbiota to elevated CO₂ in an artificial tropical ecosystems. *Journal of Microbiological Methods*, **36**, 45–54.
- Jones DL, Hodge A, Kuzyakov Y (2004) Plant and mycorrhizal regulation of rhizodeposition. *New Phytologist*, **163**, 459–480.
- Jossi M, Fromin N, Tarnawski S, Kohler F, Gillet F, Aragno M, Hamelin J (2006) How elevated pCO₂ modifies total and metabolically active bacterial communities in the rhizosphere of two perennial grasses grown under field conditions. *FEMS Microbiology Ecology*, **55**, 339–350.
- Kandeler E, Mosier AR, Morgan JA, Milchunas DG, King JY, Rudolph S, Tschirko D (2008) Transient elevation of carbon dioxide modifies the microbial community composition in a semi-arid grassland. *Soil Biology and Biochemistry*, **40**, 162–171.
- Kanerva T, Palojarvi A, Rämö K, Manninen S (2008) Changes in soil microbial community structure under elevated tropospheric O₃ and CO₂. *Soil Biology and Biochemistry*, **40**, 2502–2510.
- Kuzyakov Y, Domanski G (2000) Carbon input by plants into the soil. Review. *Journal of Plant Nutrition and Soil Science*, **163**, 421–431.
- Kuzyakov Y, Ehrensberger H, Stahr K (2001) Carbon partitioning and below-ground translocation by *Lolium perenne*. *Soil Biology and Biochemistry*, **33**, 61–74.
- Kuzyakov Y, Siniakina SV, Rühlmann J, Domanski G, Stahr K (2002) Effect of nitrogen fertilisation on below-ground carbon allocation in lettuce. *Journal of the Science of Food and Agriculture*, **82**, 1432–1441.

- Lagomarsino A, De Angelis P, Moscatelli M, Grego S (2009) The influence of temperature and labile C substrates on heterotrophic respiration in response to elevated CO₂ and nitrogen fertilization. *Plant and Soil*, **317**, 223–234.
- Lagomarsino A, Knapp B, Moscatelli M, De Angelis P, Grego S, Insam H (2007) Structural and functional diversity of soil microbes is affected by elevated CO₂ and N addition in a poplar plantation. *Journal of Soils and Sediments*, **7**, 399–405.
- Lesaulnier C, Papamichail D, Mccorkle S *et al.* (2008) Elevated atmospheric CO₂ affects soil microbial diversity associated with trembling aspen. *Environmental Microbiology*, **10**, 926–941.
- Lichter J, Barron SH, Finzi AC, Irving KF, Roberts MT, Stemmler EA, Schlesinger WH (2005) Soil carbon sequestration and turnover in a pine forest after six years of atmospheric CO₂ enrichment. *Ecology*, **86**, 1835–1847.
- Lipson D, Blair M, Barron-Gafford G, Grieve K, Murthy R (2006) Relationships Between Microbial Community Structure and Soil Processes Under Elevated Atmospheric Carbon Dioxide. *Microbial Ecology*, **51**, 302–314.
- Lipson D, Monson R, Schmidt S, Weintraub M (2009) The Trade-Off Between Growth Rate and Yield in Microbial Communities and the Consequences for Under-Snow Soil Respiration in a High Elevation Coniferous Forest. *Biogeochemistry*, **95**, 23–35.
- Mason TG, Slater JH (1979) Competition between an *Escherichia coli* tyrosine auxotroph and a prototrophic revertant in glucose- and tyrosine-limited chemostats. *Antonie van Leeuwenhoek*, **45**, 253–263.
- Mayr C, Miller M, Insam H (1999) Elevated CO₂ alters community-level physiological profiles and enzyme activities in alpine grassland. *Journal of Microbiological Methods*, **36**, 35–43.
- Monson RK, Lipson DL, Burns SP, Turnipseed AA, Delany AC, Williams MW, Schmidt SK (2006) Winter forest soil respiration controlled by climate and microbial community composition. *Nature*, **439**, 711–714.
- Montealegre CM, van Kessel C, Russele MP, Sadowsky MJ (2002) Changes in microbial activity and composition in a pasture ecosystem exposed to elevated atmospheric carbon dioxide. *Plant and Soil*, **243**, 197–207.
- Panikov NS (1995) *Microbial Growth Kinetics*. Chapman & Hall, London, 378 pp.
- Panikov NS, Sizova MV (1996) A kinetic method for estimating the biomass of microbial functional groups in soil. *Journal of Microbiological Methods*, **24**, 219–230.
- Paterson E, Thornton B, Midwood AJ, Osborne SM, Sim A, Millard P (2008) Atmospheric CO₂ enrichment and nutrient additions to planted soil increase mineralisation of soil organic matter, but do not alter microbial utilisation of plant- and soil C-sources. *Soil Biology and Biochemistry*, **40**, 2434–2440.
- Pregitzer KS, Zak DR, Maziasz J, DeForest J, Curtis PS, Lussenhop J (2000) Fine root growth, mortality, and morphology in a factorial elevated atmospheric CO₂ × soil N availability experiment. *Ecological Applications*, **10**, 18–33.
- Rillig MC, Scow KM, Klironomos JN, Allen MF (1997) Microbial carbon-substrate utilization in the rhizosphere of *Gutierrezia sarothrae* grown in elevated atmospheric carbon dioxide. *Soil Biology and Biochemistry*, **29**, 1387–1394.
- Ross DJ, Tate KR, Newton PCD (1995) Elevated CO₂ and temperature effects on soil carbon and nitrogen cycling in ryegrass/white clover turves of an Endoaquept soil. *Plant and Soil*, **176**, 37–49.
- Schlesinger WH, Lichter J (2001) Limited carbon storage in soil and litter of experimental forest plots under increased atmospheric CO₂. *Nature*, **411**, 466–469.
- Schortemeyer M, Hartwig UA, Hendrey GR, Sadowsky MJ (1996) Microbial community changes in the rhizospheres of white clover and perennial ryegrass exposed to free air carbon dioxide enrichment (FACE). *Soil Biology and Biochemistry*, **28**, 1717–1724.
- Smith JL, Mcneal BL, Cheng HH, Campbell GS (1986) Calculations of microbial maintenance rates and net mineralization in soil at steady-state. *Soil Science Society of America Journal*, **50**, 332–338.
- Stenström J, Svensson K, Johansson M (2001) Reversible transition between active and dormant microbial states in soil. *FEMS Microbiology Ecology*, **36**, 93–104.
- Taneva L, Gonzalez-Meler MA (2008) Decomposition kinetics of soil carbon of different age from a forest exposed to 8 years of elevated atmospheric CO₂ concentration. *Soil Biology and Biochemistry*, **40**, 2670–2677.
- Torbert HA, Johnson HB (2001) Soil of the intensive agriculture biome of Biosphere2. *Journal of Soil and Water Conservation*, **56**, 4–11.
- Van Groenigen K-J, Six J, Harris D, Blum H, Van Kessel C (2003) Soil ¹³C–¹⁵N dynamics in an N₂-fixing clover system under long-term exposure to elevated atmospheric CO₂. *Global Change Biology*, **9**, 1751–1762.
- van Veen JA, Liljeroth E, Lekkerkerk LJA, van de Geijn SC (1991) Carbon fluxes in plant–soil systems at elevated atmospheric CO₂ levels. *Ecological Applications*, **1**, 175–181.
- Weigel HJ, Dämmgen U (2000) The Braunschweig carbon project: atmospheric flux monitoring and free air carbon dioxide enrichment (FACE). *Journal of Applied Botany*, **74**, 55–60.
- Weigel HJ, Manderscheid R, Burkart S, Pacholski A, Waloszczyk K, Frühauf C, Heinemeyer O (2006) Responses of an Arable Crop Rotation System to Elevated [CO₂]. In: *Managed Ecosystems and CO₂. Case Studies, Processes, and Perspectives*, (eds Nösberger J, Long SP, Norby RJ, Stitt M, Hendrey GR, Blum H), pp. 121–138. Springer, Amsterdam.
- Wiesenberg GLB, Schwarzbaauer J, Schmidt MWI, Schwark L (2008) Plant and soil lipid modification under elevated atmospheric CO₂ conditions: II. Stable carbon isotopic values (δ¹³C) and turnover. *Organic Geochemistry*, **39**, 103–117.
- WRB (1998) *World Reference Base for Soil Resources*. Food and Agriculture Organization of the United Nations, Rome.
- Yadav AS (2007) Auxotrophy in rhizobia revisited. *Indian Journal of Microbiology*, **47**, 279–288.
- Zabel B, Hawes P, Hewitt S, Marino BDV (1999) Construction and engineering of a created environment: overview of the Biosphere 2 closed system. *Ecological Engineering*, **13**, 43–63.
- Zak DR, Pregitzer KS, Curtis PS, Holmes WE (2000a) Atmospheric CO₂ and the composition and function of soil microbial communities. *Ecological Applications*, **10**, 47–59.
- Zak DR, Pregitzer KS, King JS, Holmes WE (2000b) Elevated atmospheric CO₂, fine roots and the response of soil microorganisms: a review and hypothesis. *New Phytologist*, **147**, 201–222.